



NATIONAL OPEN UNIVERSITY OF NIGERIA

SCHOOL OF SCIENCE AND TECHNOLOGY

COURSE CODE: CHM 407

COURSE TITLE: REACTION KINETICS



NATIONAL OPEN UNIVERSITY OF NIGERIA

National Open University of Nigeria
Headquarters
14/16 Ahmadu Bello Way
Victoria Island
Lagos

Abuja Annex
245 Samuel Adesujo Ademulegun Street
Central Business District
Opposite Arewa Suites
Abuja

CHM 407 – REACTION KINETICS

Dr H.D. Aliyu
CHEMISTRY DEPARTMENT
UNIVERSITY OF ABUJA

TABLE OF CONTENTS	PAGE
Introduction.....	2
What you will learn in this Course.....	3
Course Aims.....	3
Course Objectives.....	3-4
Working through this Course.....	4
Course Materials	4
Study Units	5-6
Textbooks and References.....	6

Introduction

The outcome of any chemical process is determined by the thermodynamics and kinetics of the reactions involved. The study of the thermodynamics reveals whether or not a given process can occur spontaneously and the process of reactions when equilibrium is established. This will be subject to CHM 301. Reaction kinetics is concerned with the speed at which chemical changes take place. Spontaneity and speed must be favourable to observe the formation of products of chemical reactions.

This course CHM 407 concerns itself with the speed, or rates of chemical reactions. The study of reaction rates allows for the prediction of how fast a reaction mixture moves to equilibrium and how the reaction rate would be optimized by controlling certain factors such as temperature, pressure and the presence of a catalyst. The study of rate often reveals the sequence of elementary steps that lead to the product formation from the reactants, which is referred to as the reaction mechanism.

Many reactions of technologies and biological importance involve complex sequence of steps, or proceed at a useful rate only in the presence of catalysts

Many gas reactions of industrial importance occur only on the surface of solids acting as catalysts. The mode of action of such surface reactions are discussed in this course. Light is a form of energy and light energy can be used to initiate or sustain chemical reaction. This is also discussed in this course.

What you will learn in this course

The course is core for students of the Bachelor's degree in Chemistry. There are 5 units in the course. The course introduces the students to the fundamental concepts in kinetics studies and the different methods of rate studies. Methods of analyzing experimental results are discussed and applied to results from actual studies. Laws and theories are discussed and applied to account for experimental observations in rate studies. The effects of a catalyst on reaction rate are discussed and explained qualitatively and quantitatively.

The discussion of complex reactions, and the analysis of results from experiments. Physisorption and chemisorption processes are used to explain surface reactions. Finally, the study photochemistry reveals the use of light energy to initiate and sustain reactions. Light emission and absorption phenomena are also discussed and explained in the course.

A good mathematical background is required for proper understanding of the course.

Course Aims

The course aims to introduce students to the fundamental concepts in rate and mechanism studies to explain natural phenomena of technological and biological importance.

Course Objectives

We expect that after studying this course, you should be able to:

- Define rate law, rate constant and order of reaction,
- Differential between order of reaction and stoichiometry,
- State the experimental methods for studying the reaction rate,
- Derive integrated rate laws for first order, second order and zeroth order reactions and use them for calculating rate constants,
- State the methods for determining the order of reaction,
- Define elementary reaction and molecularity,
- Explain the modes of action of heterogeneous and homogeneous catalysts.
- Account for acid-base catalysis using the Bronsted theory,

COURSE GUIDE**CHM 407**

- Explain the significance of the Michael's county in enzyme catalysis.
- Explain the terms, concurrent, reversible and consecutive reactions
- Analysis experimental results for complex reactions i.e. how product concentration or reactant concentration varies with time of reaction.

- Define and use the Langumair absorption isotherm
- Explain order for a surface reaction
- State Beer-Lambers Law
- Calculate quantum yield
- List some examples of photochemical reactions
- Derive rate expressions for some photochemical reactions
- List some applications of photochemical studies.

Working Through This Course

This course involves that you would be required to spend lot of time to read. The content of this material is very dense and require you spending great time to study it. This accounts for the great effort put into its development in the attempt to make it very readable and comprehensible. Nevertheless, the effort required of you is still tremendous. I would advice that you avail yourself the opportunity of attending the tutorial sessions where you would have the opportunity of comparing knowledge with your peers.

Course Materials

You will be provided with the following material;

1. Course Guide
2. Study Units.

In addition, the course comes with a list of recommended textbooks which though are not compulsory for you to acquire or indeed read, are necessary as supplements to the course material.

Study Units

Unit 1 **Reaction rates:** Fundamental methods for rate studies; First, second, reactions; Analysis of experimental results for reaction order (integral differential and half-life methods);

Unit 2 Zeroth, third and Fast reactions; Complex reactions: concurrent, reversible and consecutive reactions and analysis of experimental results for such reactions and enzyme substrate complex and the Michael's constant.

Unit 3 **Reaction meachanism:** Uni and bimolecular reactions, chain reactions, Theories of reaction rates (Arrhenius, collision and transition state theory), Linderman hypothesis and the steady state approximation.

Unit 4 Heterogeneous reactions and catalysis: Physorption and chemisorption, Types of reaction isotherms. Orders of heterogeneous reactions. Catalyst, Heterogeneous catalysis examples and mechanism

Unit 5 Photochemical reactions: Laws of photochemistry, Quantum efficiency Photo-chemical reactions and photo-physical processes, Photosensitisation, Applications of photochemistry and chemiluminescence.

In unit 1 we discuss the different types of complex reactions namely; concurrent, reversible and consecutive (chain) reactions. The analysis of experimental results for such reaction are also explained in the unit.

Unit 2 deals with the zeroth, third and fast reaction and determination of reaction order

In Unit 3, the mechanism of uni , bimolecular reactions and chain reactions are discussed. Theories to harmonize observed rates and mechanism are discussed.

In unit 4 we introduce surface chemistry and catalysis. Physorption and chemisorption processes are discussed and the different types of reaction isotherms explained. The mode of action of the heterogeneous catalyst is further explained and examples and applications of heterogeneous catalysts discussed.

In unit 5, we discuss, light induced reactions (photochemistry). In photochemistry, we study the absorption and emission of light by matter and in photochemical reactions, reactions acquire the necessary activation energy through light absorption.

Textbooks and References

Principles of Physical chemistry, Samuel H. Maron and Carl F. Prutton, Oxford and IBH Publishing Co., New Delhi, 4th ed., 1985. 1997.

Physical Chemistry, Gilbert W. Castellan, Narosa Publishing House, Addison-Wesley/Narosa (Indian student Edition), Delhi 3rd ed., 1983.

Physical Chemistry, P.C. Rakshit, Sarat Book House, Calcutta, 5th ed., 1988.

Physical Chemistry – Principles and Problems, D.V.S Jain and S.P. Jauhar, Tata McGraw-Hill Pub. Company Ltd., New Delhi, 1988.

Physical chemistry Joseph H Niggle 3rd edition Harper Collins college publisher 1936

Introduction to physical chemistry Mark ladd 3rd edition Cambridge university press 1999

Physical chemistry Atkins PW 3rd edition oxford university press 1986

Advanced chemistry (physical and industrial) Philip Mathew Cambridge university press 2003

UNIT 1 CHEMICAL KINETICS

Structure

- 1.1 Introduction
Objectives
- 1.2 Some Fundamental Concepts
- 1.3 Experimental Method of Rate Studies
- 1.4 First Order and Second Order Reactions
- 1.5 Summary
- 1.6 Terminal questions
- 1.7 Answer

1.1 INTRODUCTION

Chemical kinetics is the study of rates and mechanisms of chemical reactions. The rate of a reaction depends on many factors such as the concentration of the reactants, temperature, Catalyst, etc

We shall start the unit explaining the dependence of rate of reaction on the concentrations of the reactants. In this process, we shall be defining the terms such as rate equation, rate law and order of a reaction. We shall then derive first order, second order of reaction.

Objectives

After studying this unit, you should be able to:

- define rate law, rate constant and order of reaction,
- differentiate between order of reaction and stoichiometry
- state the experimental methods for studying the reaction rates,
- derive integrated rate laws for first order, second order reactions and use them for calculating rate constants,

Throughout this unit, the phrase 'rate of reaction' means instantaneous reaction rate.

We shall shortly study the method of arriving at the reaction rates from the values of concentrations of a component at different time intervals.

The symbol ' Δ ' is to be read as delta. It denotes change in a property.

Convention For Expressing Reaction Rates

In order to obtain a single value for the reaction rate, it is necessary to divide the rate of consumption of a reactant or the rate of formation of a product by the stoichiometric coefficient of the respective species. To illustrate this, let us consider the reaction:



In this reaction, A and B are reactants, and C and D are products; a , b , c and d are the stoichiometric coefficients. The reaction rate is related to the rates of consumption of the reactants and the rates of formation of the products as follows:

Reaction rate

$$= \frac{1}{a} \frac{-d[A]}{dt} = \frac{1}{b} \frac{-d[B]}{dt}$$

$$= \frac{1}{c} \frac{d[C]}{dt} = \frac{1}{d} \frac{d[D]}{dt}$$

... (18.2)

Using this general equation, you examine E.q 1.1

The number preceding the formula of a substance in the balanced equation is its stoichiometric coefficient.

1.2 SOME FUNDAMENTAL CONCEPTS

In this section, we shall define some terms such as rate of Reaction, rate law, rate constant, order of a reaction and stoichiometry.

Rate of Reaction

The rate of reaction or the velocity of reaction at a specified time is defined as the decrease in the concentration of a reactant or the increase in the concentration of a product per unit time. The rate of reaction at a specified time is also known as instantaneous rate of reaction; it can be generally defined as the rate of change of concentration of a specified species at a particular instant. While specifying the reaction rate, we must mention the component with respect to which it is stated. Let us consider a simple reaction,



As per the reaction stoichiometry, one molecule of B is formed for every molecule of A consumed. The reaction rate can be specified in the following ways:

- We can measure the concentration of the reactant A at various time intervals. From these values, we can specify the decrease in concentration of A with respect to time at any particular instant. The reaction rate thus obtained is the rate of consumption of A.

$$\text{Rate of consumption of A} = \frac{\text{Decrease in the concentration of A}}{\text{Change in time}}$$

$$= \frac{-\Delta[A]}{\Delta t}$$

- $\Delta [A]$ means change in the concentration of A and $-\Delta [A]$ means the decrease in the concentration of A. While writing a rate expression with respect to a reactant, there is a preceding negative sign (since it is customary to express the rate of a reaction as a positive quantity).
- We can measure the concentration of the product B at various time intervals. From these values, we can arrive at the rate of formation of B at any particular instant.

$$\text{Rate of formation of B} = \frac{\text{Increase in the concentration of B}}{\text{Change in time}}$$

$$= \frac{\Delta[B]}{\Delta t}$$

The rates of consumption of reactants and the rates of formation of products are related through their stoichiometric coefficients. For example, consider the decomposition of NO_2 .



We can write the relationship between the rates of consumption of NO_2 and the rates of formation of NO and O_2 as follows:

$$\begin{aligned} \frac{1}{2} (\text{Rate of; consumption of } \text{NO}_2) &= \frac{1}{2} (\text{Rate of formation of } \text{NO}) \\ &= \text{Rate of formation of } \text{O}_2 \end{aligned} \quad \dots (1.1)$$

Using the convention as expressed by Eq. 1.2, we can write as shown below;

$$\text{Reaction rate} = \frac{1}{2} \left(\frac{-d[\text{NO}_2]}{dt} \right) = \frac{1}{2} \left(\frac{d[\text{NO}]}{dt} \right) = \left(\frac{d[\text{O}_2]}{dt} \right) \dots(1.2)$$

You can understand the above relationship if you bear in mind that if two molecules of NO_2 are consumed, two molecules of NO and one molecule of O_2 are formed. In other words, the reaction rate is equal to:

- half the rate of consumption of NO_2
- half the rate of formation of NO , and
- the rate of formation of O_2

Calculation of Reaction Rate

You may be curious to know as to how the reaction rates are calculated. Take for instance the following reaction:



In Figs 1.1 a, b, and c, you can see concentration (c) against time (t) plots for NO_2 , NO and O_2 as per the values given in Table 18.1. In these figures, the graphical method of calculation of the reaction rates for the consumption of NO_2 and for the formation of NO and O_2 are illustrated. The reaction rate at any particular instant is obtained by calculating the slope of a line tangent to the curve at that point.

Table 1.1 Concentration of NO_2 and NO and O_2 at Different Time interval at 673 K.

Time/s	$[\text{NO}_2]/\text{M}$	$[\text{NO}]/\text{M}$	$[\text{O}_2]/\text{M}$
0	0.00100	0	0
50	0.0079	0.0021	0.0011
100	0.0065	0.0035	0.0018
150	0.0055	0.0045	0.0023
200	0.0048	0.0052	0.0026
250	0.0043	0.0057	0.0029
300	0.0038	0.0062	0.0031
350	0.0034	0.0066	0.0033




Fig. 1.1 a) Concentration against time plot for NO_2 ; note the falling nature of the curve which is characteristic of concentration against time plot for a

From the slope of the tangent time drawn (corresponding to a particular time) to the concentration (c) against time (t) curve for a component, we can obtain the rate of the reaction.

Rate of reaction

$-\text{(Slope of tangent to the } c \text{ against } t \text{ curve for the reactant)}$
 Stoichiometric coefficient of the reactant

$\text{Slope of tangent to the } c \text{ against } t \text{ curve for the product}$
 Stoichiometric coefficient of the product

The concentration of components are given in molarity (M) unit $1 \text{ M} = 1 \text{ mol dm}^{-3}$

reactant.

Rate of consumption of NO_2 ; $= - \text{Slope of the tangent line at } t = 200 \text{ s}$

at $t = 200 \text{ s}$

$$\begin{aligned} &= -(-1.31 \times 10^{-5}) \text{ Ms}^{-1} \\ &= 1.31 \times 10^{-5} \text{ Ms}^{-1} \end{aligned}$$

$$\begin{aligned} \text{Reaction rate} &= \frac{1}{2} (\text{Rate of consumption of NO}_2) = \frac{1}{2} 1.31 \times 10^{-5} \text{ Ms}^{-1} \\ &= 6.55 \times 10^{-6} \text{ Ms}^{-1} \end{aligned}$$

- b) concentration against time plot for NO; note the rising nature of the curve which is characteristic of concentration against time plot for a product.

$$\begin{aligned} \text{Rate of formation of NO} &= \text{Slope of the tangent line at } t = 200 \text{ s} \\ \text{at } t = 200 \text{ s} &= 1.30 \times 10^{-5} \text{ Ms}^{-1} \end{aligned}$$

$$\begin{aligned} \text{Reaction rate} &= \frac{1}{2} (\text{Rate of formation of NO}) \\ &= 1/2 \times 1.30 \times 10^{-5} \text{ M s}^{-1} \\ &= 6.50 \times 10^{-6} \text{ M s}^{-1} \end{aligned}$$

- c) concentration against time plot for O₂; note again the rising curve. Compared to the curve for NO, the curve for O₂ rises slowly.

$$\begin{aligned} \text{Rate of formation of O}_2 &= \text{Slope of the tangent line at } t = 200 \text{ s} \\ \text{at } t = 200 \text{ s} &= 6.25 \times 10^{-4} \text{ Ms}^{-1} \end{aligned}$$

$$\begin{aligned} \text{Reaction rate} &= \text{Rate of formation of O}_2 \\ &= 6.25 \times 10^{-6} \text{ Ms}^{-1} \quad \text{i} \end{aligned}$$

For the curves (a) and (b), the tangents are not indicated.

From the slope values at $t = 200 \text{ s}$, you can see that the following relationship is nearly correct.


$$\begin{aligned} \text{Reaction rate} &= \frac{1}{2} (\text{Rate of consumption NO}_2) = \frac{1}{2} (\text{Rate of formation of NO}) \\ &= \text{Rate of formation of O}_2 \end{aligned}$$

In our discussion, we are mainly interested in the concentration against time plots for the reactants. In other words, we want to study the reactions under conditions where the rate of the forward reaction is significant but the reverse reaction rate is low. This is made possible, if we study the reaction upto a point where the product amounts are not high. For example, in the decomposition of NO₂, there could be a decrease in the concentration of NO₂ upto a particular time. Afterwards, enough nitric oxide and oxygen are formed and the reverse reaction also could take place leading to the formation of NO₂. In order to simplify the

situation, it is better to study the reaction rates before significant amounts of products are formed. In general, the rates of reactions are complex functions of the concentrations of the reactants and the products at a given temperature. However, there are some reactions in which the rates are proportional to the simple powers of the concentrations of the reactants. We shall be mostly concerned with this class of reactions.

Decomposition of N₂O₅

The decomposition of N₂O₅ in the gas phase was studied at 323 K



The instantaneous rates of this reaction calculated from [N₂O₅] against time plot (similar to Fig. 1.1) are given in Table 1.2

Table 1.2: Rates for the Decomposition of N₂O₅ at 323 K

[N ₂ O ₅]/M	Rate/MS ⁻¹	Rate / [N ₂ O ₅]
(i) 0.300	(ii) 2.73 × 10 ⁻⁴	(iii) 9.1 × 10 ⁻⁴
0.150	1.37 × 10 ⁻⁴	9.1 × 10 ⁻⁴
0.100	9.10 × 10 ⁻³	9.1 × 10 ⁻⁴

From columns (i) and (ii), you can see that the rate for the decomposition of N₂O₅ decreases with the decrease in the concentration of N₂O₅. Further, column (iii) gives the ratio of the rate to the concentration of N₂O₅. In all the three cases, it is a constant. This shows that the rate is directly proportional to the concentration of N₂O₅


$$\text{i.e. } \frac{\text{Rate}}{[\text{N}_2\text{O}_5]} = k \quad \dots (1.3)$$

$$\text{Hence, rate} = k [\text{N}_2\text{O}_5] \quad \dots (1.4)$$

where *k* is proportionality constant.

Decomposition of Hydrogen Iodide

The decomposition of hydrogen iodide was followed at a constant temperature



The instantaneous rates of this reaction were calculated using the [HI] against time plot like that of Fig. 1.1. These values are given in Table 1.3

Table 1.3: Rates for the Decomposition of HI

[HI]/M	Rate/Ms-1	Rate [HI]	Rate _{Ms} [HI] ²
(i)	(ii)	(iii)	(iv)
3.00×10^{-2}	3.60×10^{-5}	1.2×10^{-3}	4.00×10^{-2}
2.00×10^{-2}	1.60×10^{-3}	8.0×10^{-4}	4.00×10^{-2}
1.50×10^{-2}	9.01×10^{-6}	6.0×10^{-4}	4.00×10^{-2}

From Table 1.3, you can see that the rate of decomposition of HI decreases with decrease in the concentration of HI, as in the case of the decomposition of N₂O₅. Further, it is evident from column (iii) that rate/[HI] is not a constant. But, as per column (iv), rate/[HI]² is a constant.

From Table 18.3, it is evident that

$$\text{Rate} / [\text{HI}]^2 = k \quad \dots(1.5)$$

$$\text{Hence, rate} = k [\text{HI}]^2 \quad \dots (1.6)$$

where k is a proportionality constant

For many chemical reactions, the relationship between the reaction rate and the concentration can be expressed in a simple way as in Eq. 1.4 or 1.6. We shall first consider these simple cases. Later we shall discuss those reactions for which the rate concentration relationship is more complex,

Rate Law Rate Constant

The relationship expressed as in Eq. 1.4 or Eq. 1.6 is called the **rate law**. A rate law is an equation expressing the relationship between the instantaneous reaction rate and the concentrations of the reactants in a reaction.

The rate law for a simple reaction with one reactant may be of the following type:

$$\text{Reaction rate} = k [\text{Reactant}]^n \quad \dots(1.7)$$

where k is called the rate constant or rate coefficient or the specific rate or the reaction.

Thus by definition, the rate constant is independent of concentration, but it may depend on other factors. In this equation, n refers to the order of the reaction. The

order with respect to a component is the power to which the concentration of that component is raised in the rate law. Comparing Eq. 1.7 with Eqs 1.4 and 1.6 we conclude that

- i) $n = 1$ in Eq. 1.4; i.e.; decomposition of N_2O_5 is a **first order** reaction. The significance of this statement is that the reaction rate is proportional to the first power of concentration of N_2O_5 .

$$\text{i.e., Rate} = k [N_2O_5]^1 \quad \dots (1.8)$$

where k is the **first order rate constant**.

From Eq. 18.7, it can be seen that if $[\text{reactant}] = 1$ then $k = \text{rate}$. For this reason, k is called the specific rate.

- ii) $n = 2$ for the decomposition of HI; i.e., the decomposition of HI is a **second order reaction**. Again this means that the decomposition rate of HI is proportional to the second power or square of the concentration of HI.

$$\text{i.e., Rate} = k [HI]^2 \quad \dots(1.9)$$

where k is the **second order rate constant**.

Order of Reaction and Stoichiometry

The rate laws as well as the order of the reaction must be determined experimentally; these cannot be predicted from the stoichiometry of the reaction. The stoichiometry of reaction gives the relationship between the amounts of the reactants and the amount of the product. The stoichiometry of a reaction must be differentiated from the order of a reaction. Let us consider the following examples,

Example 1

The gas-phase decomposition N_2O_5 yields NO_2 and O_2 at a particular temperature.



The experimentally observes rate law for the reaction rate $= k [N_2O_5]$
Comment on the order and the stoichiometry of the reaction.

Solution

It can be seen that the stoichiometric coefficient of N_2O_5 is 2 whereas the order of reaction is 1.

Example 2

The balanced equation for the decomposition of nitrous oxide is given below:



The rate law is,


$$\text{rate} = k [\text{N}_2\text{O}_5]$$

Comment on the order of the reaction and the stoichiometry.

Solution

Again the stoichiometric coefficient of N_2O_5 is 2 whereas the order of reaction is 1.

In the above two examples, the order of reaction and the stoichiometry are not identical, but there are cases where: the order and stoichiometric coefficient are identical. One of the examples can be seen in the following reaction:



$$\text{Rate} = k [\text{HI}]^2$$

In the decomposition of HI, the order of reaction is two. The stoichiometric coefficient of HI is also 2.


From the above examples, you can see that the stoichiometric coefficient and the order of the reaction need not be the same always. You must bear in mind the following points while arriving at a rate law.

- i) In the case of simple reactions, the Concentrations of the reactants appear in rate law; but the Concentrations of the products do not appear in the rate law. It is so since the rate measurements are done under the conditions where the reverse reaction rate is negligibly low.
- ii) The order of the reaction must be determined experimentally; the experimental methods will be discussed in the next section.
- iii) The order of a Reaction need not be identical with the stoichiometric coefficient of the reactant.

So far we considered the reactions involving only one reactant. In case of reactions involving many reactants, the rate of a reaction may depend on the concentrations of more than one reactant. In such cases, we can calculate the order of the reaction with respect to the individual reactant and also the **overall**

order. The overall order is the sum of the powers to which the individual concentrations are raised in the rate law.

In general, for a reaction,




if the rate law is experimentally found to be,

$$\text{rate} = k [A]^m [B]^n [C]^p$$

then, the overall order of the reaction = $m + n + p$.

For example in the following reaction,



The overall rate of the reaction is four, being first order in BrO_3^- , first order in Br^- second order in H^+

The rate laws discussed so far are called **differential rate laws**. Such rate laws describe the dependence of reaction rate on concentration. From these differential rate laws, we can obtain the **integrated rate laws** through integration. These integrated rate laws help us in relating the concentration of a substance to time. In order words, using the integrated rate laws, we can calculate the concentration of a substance at any specific time. In Secs. 18.4 and 18.5, we shall discuss the derivation of the integrated forms of rate laws. In the next section, we shall discuss some experimental methods of studying the reaction rates.

SAQ1

At 323 K, the rate of reaction for the decomposition of N_2O_5 at a particular instant is $2.74 \times 10^{-4} \text{ M s}^{-1}$. Calculate the rate of formation of O_2 . The reaction is represented below:



.....
.....
.....

SAQ2

In the decomposition of hydrogen iodide, what is the relationship between the rate of decomposition of HI and the rate of formation of H_2 ?

.....
.....
.....

For the $\text{BrO}_3^- - \text{Br}^- - \text{H}^+$ reaction let us compare the order of reaction and the stoichiometric coefficient for each reactant.

	BrO_3^-	Br^-	H^+
Order	1	1	1
Stoichiometric Coefficient	1	5	6

It may be noticed that the stoichiometric coefficients and the respective orders of reaction are not identical throughout.

1.3 EXPERIMENTAL METHODS OF RATE STUDIES

Many physical and chemical methods are available for studying the reaction rates. Some of them are listed below:

i) Volume or Pressure Measurement

When one or more of the components are gases, the reaction rate can be followed by measuring the volume or pressure change. The partial pressures of the species are to be calculated using the reaction stoichiometry.

Spectrophotometers have arrangements for generation of nearly monochromatic radiation in visible and ultraviolet regions and also for the measurement of radiation transmitted by the absorbing substance.

Nowadays many sophisticated instruments such as nuclear magnetic resonance spectrometer, mass spectrometer etc, are used in reaction kinetics, We shall not discuss these aspects.

ii) Titrimetry

Using acid-base or oxidation-reduction titrations, the reaction course can be followed if at least one of the components in the reaction is an acid or a base or an oxidising agent or a reducing agent.

iii) **Conductometry or Potentiometry**

If one or more of the ions are present or produced in the reaction, suitable methods can be designed based on conductivity or potentiometric measurements.

iv) **Spectrophotometry**

When a component of the reaction has a strong absorption band at a particular wavelength region, spectrophotometers could be used for measuring the reaction rate. Photoelectric colorimeters are cheaper instruments and are mainly useful for reaction; rate studies in visible region.

v) **Polarimetry**

When at least one of the components of a reaction is optically active, the reaction rate can be studied from the measurements of optical rotation:


Depending of the reaction under study, the concentration of a reactant or a product is followed at various time intervals using any of the methods mentioned above. These values are then used for calculating the rate constant. Examples are worked out in the next section to illustrate the rate constant calculation. Before studying these examples, we shall discuss the method of arriving at the integrated forms of rate laws.

SAQ 3

State the name of a suitable, experimental method that can be followed to monitor the reaction rate in each following cases.



Hint: Bromine absorbs strongly in the visible region, while hydrogen and hydrogen bromide do not.



1.4 FIRST ORDER AND SECOND ORDER REACTION

In this section, we shall derive integrated rate laws for first and second order reactions.

Integrated Rate Law for First Order Reactions.

Let us consider the following reaction, which is experimentally found to be first order



Applying Eq. 1.7 we can write

$$\text{rate} = \frac{-d[A]}{dt} = k[A]^1 \quad \dots(1.13)$$

where k is the first order rate constant and $n = 1$. This means that the rate of consumption of A at any given time is directly proportional to the first power of the concentration of A at that time.

In order to obtain the integrated rate for first order reaction concentrations of A at the start of the reaction and at a time t as mentioned below:

At time = 0 (i.e., at the start), the concentration of A = $[A]_0$

At time = t the concentration of A = $[A]_t$

Using these limits of concentration and time, we can integrate Eq. 1.13 after rearranging it as follows:

$$\int_{[A]_0}^{[A]_t} - \frac{d[A]}{[A]} = \int_0^t k dt \quad \dots (1.14)$$

$$\text{i.e., } - \{ \ln [A] \} = k[t] \int_0^t \frac{[A]_t}{[A]_0} \quad \dots (1.15)$$

$$- \{ \ln [A]_t - \ln [A]_0 \} = k (t - 0) \quad \dots (1.16)$$

$$\ln [A]_0 - \ln [A]_t = kt \quad \dots(1.17)$$

$$\text{Hence, } \ln \frac{[A]_0}{[A]_t} = kt \quad \dots(1.18)$$

Eq. 1.18 can be written in the exponential form as follows:

$$[A]_t = [A]_0 e^{-kt} \quad \dots(1.19)$$

In order to understand the transformation of Eq. into Eq 1.16.

The transformation of Eq. 1.18 into Eq. 1.19 could be explained as follows

$$\ln \frac{[A]_0}{[A]_t} = kt \quad 1.18$$

$$-\ln \frac{[A]_t}{[A]_0} = kt$$

Taking antilogarithms

$$\frac{[A]_t}{[A]_0} = e^{-kt}$$

$$[A]_t = [A]_0 e^{-kt} \quad 1.19$$

Corresponding to Eq. 1.19, we can draw the concentration vs time plot for a first order reaction. This curve is of the type shown in Fig. 1.2. Such a curve is called an exponential decay curve. In an exponential decay curve, there is a steep decrease in concentration initially. It is followed by a slow decrease in concentration subsequently. Note that the decay curve goes parallel to x -axis after longer time intervals indicating that the reaction will take infinite time for completion.




Fig 1.2: Exponential decay of A.

Radioactive decay is a good example of first order/reaction. A detailed discussion on radioactive decay has been given in CHE 213

Let us come back to Eq. 1.18 It is convenient to work with logarithms to the base 10 (known as common logarithms) Hence we can rewrite Eq. 1.18 as follows:

Remember the following formula: $\ln x = 2.303 \log x$

$$2.303 \log \frac{[A]_0}{[A]_t} = kt$$

... (1.20)

$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]_t}$$

Eq. 1.20 is useful in calculating the concentration of the reactant at a time t (i.e., $[A]_t$) provided, its initial concentration (i.e., $[A]_0$), k and t are known. Also, k can be calculated if $[A]_0$, $[A]_t$ and t are known. Using the graphical method, we can test whether a reactant follows first order or not.

Graphical Method of Calculating First Order Rate Constant

In order to facilitate graphical representation, Eq. 18.20 is modified as follows

$$\log \frac{[A]_0}{[A]_t} = \frac{kt}{2.303} \quad \dots (1.21)$$

$$\text{Hence } \log [A]_0 - \log [A]_t = \frac{kt}{2.303} \quad \dots (1.22)$$

$$\text{i.e., } -\log [A]_t = \frac{kt}{2.303} - \log [A]_0 \quad \dots (1.23)$$

$$\log [A]_t = \frac{-k}{2.303} t + \log [A]_0 \quad \dots (1.24)$$

From Eq 1.21 we can derive Eq. 1.22 by substituting the concentrations of A at the intervals t_1 , and t_2 as follow:

At time = t_1

concentration of A = $[A]_1$

At lime = t_2 ,

concentration of A = $[A]_2$

$$\log \frac{[A]_0}{[A]_1} = \frac{k t_1}{2.303}$$

$$\log \frac{[A]_0}{[A]_2} = \frac{k t_2}{2.303}$$

Hence

$$\begin{aligned} \log \frac{[A]_0}{[A]_2} - \log \frac{[A]_0}{[A]_1} \\ = \frac{k}{2.303} (t_2 - t_1) \quad \dots(1.26) \end{aligned}$$

$$\text{i.e } \log \frac{[A]_1}{[A]_2} = \frac{k}{2.303} (t_2 - t_1) \quad \dots(1.27)$$

By comparing Eq. 1.24 with the equation for a straight line,
 $y = mx + c$,

We can infer that by plotting $\log; [A]$, against time a straight line must be obtained for a first order reaction. Such plots are shown in Figs. 1.3 1.5. The rate constant can be calculated from the slope as follows:

$$k = -2.303 \times \text{slope} \quad \dots (1.25)$$

Knowing the concentration of a reactant undergoing first order reaction at a particular time, it is possible to calculate its concentration at another time interval using Eq. 1.27

$$\log \frac{[A]_1}{[A]_2} = \frac{kt}{2.303} (t_2 - t_1) \quad \dots (1.27)$$

The following hints may be useful while calculating the rate constants using the integrated rate law:

- i) If the concentrations of the reactant are given at various time intervals, the concentration at $t = 0$ is equal to] $[A]_0$ and the concentration at any given

time is $[A]_t$; $\log [A]_t$ against t plot is made as described earlier. This principle is followed in Example 3.

- ii) Instead of giving the concentrations of a **reactant** at various time intervals, parameters such as partial pressures, absorbances, volumes, titre values etc. which are, proportional to the concentration of the reactant may be given. In these cases, the measurements made at zero time and at any given time t may be used instead of $[A]_0$ and $[A]_t$. For example, we can substitute the partial pressure of the reactant instead of its concentration in Eq. 1.21 and obtain Eq. 1.28.

$$\log \frac{(PA)_0}{(PA)_t} = \frac{kt}{2.303}$$

where $(PA)_0$ and $(PA)_t$ are the partial pressures of the reactant at the start and after a time, t . In Example 4, we Use this method.

- iii) Sometimes the rate of measurement are Made in terms of the concentrations of the product formed. If the stoichiometry of the reaction is such is that one molecule of the product is formed when one molecule of the reactant is consumed then the concentration of the product at $t = \infty$ must be equal to initial concentration of the reactant. Let us assume that the concentration of the product at any given time is x . Then x also represents the decrease in the concentration of the reactant after a time, t . Thus $[A]_0 =$ concentration of the product at $t = \infty$ and $[A]_t = [A]_0 - x$. Using these (relationships, Eq. 1.21 could be written as,

$$\log \frac{[A]_0}{[A]_0 - x} = \frac{kt}{2.303}$$

Calculation of this type is illustrated in Example 5.

Let us work out some examples to illustrate the calculation of k by graphical method.

Example 3

On heating cyclopropane to 770 K, it is converted into propene. In one experiment, the following data were obtained:

$t/s.$	0	300	600	900
[Cyclopropane] M	1.50×10^{-3}	1.24×10^{-3}	1.00×10^{-3}	8.3×10^{-4}

Using graphical method, test whether the above data satisfy first order rate equation. Calculate the rate constant.

Solution

Using the data, $\log [\text{cyclopropane}]/M$ values are calculated and tabulated along with t values.

$\log [\text{cyclopropane}]/M$	-2.82	-2.91	-3.00	-3.08
t/s	0	300	400	900




Fig. 1.3 $\log [\text{cyclopropane}]$ against t plot


$\log [\text{cyclopropane}]$ vs. t plot is shown in Fig. 1.3.

The graph is a straight line, confirming that the reaction is first order.

$$k = -2.303 \times \text{slope} = -2.303 \times (-2.88 \times 10^{-4}) \text{ s}^{-1} \\ = -6.63 \times 10^{-4} \text{ s}^{-1}$$

Example 4

Azomethane ($\text{CH}_3 - \text{N} = \text{N} - \text{CH}_3$) decomposes at 600 k as per the equation:



The reaction rate was followed by measuring the partial pressure of azomethane (PA) at different time intervals and the data are given below:

PA/pa	10.9	7.6	5.3	3.7	2.6
t/s	0	1000	2000	3000	4000

Using the data, test whether the reaction follows first order kinetics and calculate the rate constant.

Solution

The concentration of azomethane (CA) can be related to its partial pressure (PA.) as follows:

$$P_A = C_A RT \text{ or } P_A \propto C_A$$

So we, can use partial pressure of azomethane in Eq. 1.21 instead of concentration as shown in Eq. 1.28 Rearranging Eq; 1.28 we get,

where $(P_A)_0$ and $(P_A)_t$ are the partial pressures of azomethane at the start and after a time, t . We tabulate the data accordingly:

$\log(P_A)_t/P_A$	1.04	0.88	0.72	0.57	0.41
t/s	0	1000	2000	3000	4000




Fig. 1.4: $\log (P_A)_t$ against t plot for the decomposition of azomethane

$\log (P_A)_t$ is plotted against t as shown in Fig.1.4

From Fig. 1.4, we see that the graph is a straight line. Hence, the decomposition of azomethane is a first order reaction.

$$\begin{aligned} \text{Slope} &= 1.54 \times 10^{-4} \text{ s}^{-1} \\ k &= -2.303 \times \text{slope}, = -2.303 \times (-1.54 \times 10^{-4} \text{ S}^{-1}) \\ &= 3.55 \times 10^{-4} \text{ S}^{-1} \end{aligned}$$

Example 5

The hydrolysis rate of benzene diazonium chloride ($\text{C}_6\text{H}_5\text{N} = \text{NCl}$) in its aqueous Solution was followed by measuring the volume of nitrogen evolved at different time intervals. The reaction was found to be first order. Using the data given below, calculate the first order rate constant.

Time /s	0	1500	3000	4500	∞
$\frac{\text{Volume of N}_2}{\text{Cm}^3}$	0	6.4	12.1	17.6	81

Solution

In this example, the rate measurement is done in terms of the product. The volume (V_t) of nitrogen produced at any given time, t , is proportional to the concentration of benzene diazonium chloride reacted (x). The volume at $t \infty$ (i.e., V_∞) indicates the volume of nitrogen produced by the complete hydrolysis of benzene diazonium chloride and, it is proportional to the initial concentration of benzene diazonium chloride.

$$\text{So,} \quad [\text{A}]_0 \propto V_\infty$$

$$x \propto V_t$$

$$\text{and} \quad [\text{A}]_0 - x \propto V_\infty - V_t$$

$$\text{Hence} \quad \frac{[\text{A}]_0}{[\text{A}]_0 - x} = \frac{V_\infty}{V_\infty - V_t}$$

Using this relationship in Eq.1.29,

$$\log \frac{V_\infty}{V_\infty - V_t} = \frac{kt}{2.303}$$

$$\text{i.e.,} \quad \log (V_\infty - V_t) = \log V_\infty - \frac{kt}{2.303}$$

Using the data given above, $(V_\infty - V_t)$ and $\log(V_\infty - V_t)$ values are tabulated for different t values.

Time/s	0	1500	3000	4500
$(V_\infty - V_t)/\text{cm}^3$	81	74.6	68.9	63.4
$\log(V_\infty - V_t)/\text{cm}^3$	1.908	1.873	1.838	1.802




Fig. 1.5 $\log(v_\infty - v_t)$ against t plot for the hydrolysis of benzene diazonium chloride

In a first order reaction, the amount of the reactant remaining after n half-life periods is given by the formula:

$\log(V_\infty - V_t)$ is plotted against t as shown in Fig. 1.5

From Fig. 1.5

$$\begin{aligned} \text{slope} &= -2.36 \times 10^{-5} \text{ s}^{-1} \\ k &= -2.303 \times \text{slope} \\ &= 2.303 \times (-2.36 \times 10^{-5} \text{ s}^{-1}) \\ &= 5.44 \times 10^{-5} \text{ s}^{-1} \end{aligned}$$

In a first order reaction, the amount of the reactant remaining after n half-life periods is given by the formula:

$$\left\{ \begin{array}{l} \text{Concentration} \\ \text{of the reactant} \\ \text{remaining after} \\ n \text{ half-lives} \end{array} \right\} = \left(\frac{1}{2} \right)^n \times \left\{ \begin{array}{l} \text{Initial} \\ \text{concentration} \\ \text{of the} \\ \text{reactant} \end{array} \right\}$$

Half-Life of First Order Reactions

The time taken for the concentration of a reactant to fall to half its initial value is called the half-life of a reaction. It is denoted by the symbol, t . We can derive an expression useful in calculating the half-life of a substance undergoing first order reaction using Eq. 1.21.

$$\log \frac{[A]_0}{[A]_t} = \frac{kt}{2.303} \quad \dots(1.21)$$

We must bear in mind that when

$$t = t^{1/2}, [A]_t = [A]_0/2$$


Using these relationships in Eq. 1.21, we get,

$$\log \frac{[A]_0}{[A]_0/2} = \frac{kt_{1/2}}{2.303} \quad \dots(1.21)$$

i.e $\log 2 = \frac{kt_{1/2}}{2.303}$

or $t_{1/2} = \frac{2.303 \log 2}{k} = \frac{0.693}{k} \quad \dots(1.30)$

From Eq. 1.30; we understand that $t^{1/2}$ does not depend on initial concentration of the substance in the case of a first order reaction. This means that for a given first order reaction, half-life period is the same, whatever be the initial concentration; this leads to an interesting result that the time taken for the changes in the concentrations of the reactant such as,



is the same. The half-life periods required to bring about decrease in concentration of a reactant upto 12.5% of its initial concentration can be represented as in Fig, 1.6




Fig 1.6: Change in % $[A]_0$ over the first three half-life periods of a first order reaction) $t_{1/2}$ is the half-life period for the conversion, $[A]_0 \rightarrow 50\% [A]_0$.

Example 6

The first-order rate constant for the decomposition of N_2O_5 at 340 K is $5.20 \times 10^{-3} \text{ s}^{-1}$. Calculate the time required for the concentration of N_2O_5 to fall to (a) one-half and (b) one-fourth of its initial value.

Solution

a) Using Eq. 1.30, $t_{1/2} = \frac{0.693}{k}$

$$= \frac{0.693}{5.20 \times 10^{-3} \text{ s}^{-1}}$$

$$= 133\text{s}$$

Hence, time taken for the concentration of N_2O_5 to decrease by 50% is 133 s.

b) The time required for the decrease in the concentration of N_2O_5 to 25% of its initial value is twice the half-life period, i.e., 266 s.

Having studied the equations useful in calculating the first order rate constant half-life period of the reactant, let us derive similar equations for second order reactions.

Integrated Rate Laws for Second Order Reactions

There are two types of second order reactions.

- i) A single reactant could give rise to products through a second order reaction.




The rate equation is

$$\text{rate} = \frac{-d[A]}{dt} = k[A]^2 \quad \dots (1.33)$$

where k is the second order rate constant.

Two examples of this type are given below:






- ii) Two different reactant molecules could react to give products through a second order reaction.



The rate equation is

$$\text{reaction rate} = k [A] [B] \quad \dots(1.35)$$

Examples of this type are given below. Note that the stoichiometric ratio of the reactants is 1:1.



Again the study of these reactions could be simplified if the initial concentrations of both the reactants ($[A]_0$ and $[B]_0$) are the same, i.e.,

i.e. $[A]_0 = [B]_0$ and so $[A] = [B]$

where $[A]$ and $[B]$ are concentrations of the reactants at any given time, t

Then, Eq. 1.35 takes the same form as Eq.1.33

$$\begin{aligned} \text{Rate} &= \frac{-d[A]}{dt} = k[A][B] = k[A][A] \\ &= k[A]^2 \end{aligned} \quad \dots(1.33)$$

Thus, we could see that the rate equation takes the same form for a second order reaction, if the reaction is:

- i) second order in a single reactant or
- ii) first order in each of the two reactants such that the concentrations of the two are same throughout the reaction.

We use Eq.18.33 as the rate law for both these two types. We derive the integrated rate law for these two under case (i).

Case (i): Integrated Rate Law for a Reaction that Follows Differential Rate Law as per Eq. 1.33

We start with the differential rate law,

$$-\frac{d[A]}{dt} = k[A]^2 \quad \dots(1.33)$$

The integrated form of this equation can be obtained using the following limiting conditions:

At time = 0 (i.e., at the start), the concentration of A = [A]₀. At time = t, the concentration of A = [A]_t.

Applying these limits on the rearranged form of Eq. 1.33, we get,

$$-\int_{[A]_0}^{[A]_t} \frac{d[A]}{[A]^2} = \int_0^t k dt \quad \dots(1.36)$$

i.e., $-\left[-\frac{1}{[A]}\right]_{[A]_0}^{[A]_t} = k(t-0) \quad \dots(1.37)$

$$-\left[-\frac{1}{[A]_t} + \frac{1}{[A]_0}\right] = kt \quad \dots(1.38)$$

$$\frac{1}{[A]_t} + \frac{1}{[A]_0} = kt \quad \dots(1.39)$$

or $\frac{1}{[A]_t} = \frac{1}{[A]_0} + kt \quad \dots(1.40)$

The second order rate constant can be calculated by plotting $1/[A]_t$ against t . A straight line curve must be obtained, if the reaction is second order in the reactant. The slope of the straight line gives the second order rate constant.

$$k = \text{Slope} \quad \dots(1.41)$$

Each successive half-life is double the preceding half-life in a second order reaction. In a first order reaction, all successive half-life periods are same.

We shall later illustrate this method in Example 8.

Half-Life of a Second Order Reaction

For reactions following second order rate as per 1.33, an equation could be derived which is useful in calculating the half-life period.

At the half-life period ($t = t_{1/2}$), $[A]_t = [A]_0/2$. Using this in Eq.1.40,

$$\frac{1}{[A]_0/2} = \frac{1}{[A]_0} + kt_{1/2}$$

i.e., $\frac{2}{[A]_0} = \frac{1}{[A]_0} + kt_{1/2}$




Fig. 1.7: Change in % $[A]_0$ over three successive half-lives in a second order reaction t_1 refers to half-life for the change in concentration from $[A]_0$ to 50% $[A]_0$

$$\dots(1.42)$$

$$t_{1/2} = \frac{1}{k [A]_0}$$

From Eq.1.42, we understand that $t_{1/2}$ is inversely proportional to initial concentration for a second order reaction. As initial concentration of the reactant decreases, $t_{1/2}$ increases. If for the decrease in concentration of A to 50% of its initial value, time required is 100 s, then for the change from 50% A to 25% A, it will require 200 s.

Three successive half-lives for a second order reaction can be represented by Fig.1.7

Example 7

At 700 K, the second order rate constant for the reaction,



is $1.83 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$. Calculate the time taken for $1.00 \times 10^{-2} \text{ M}$ HI to fall to (a) one-half and (b) one-eighth of its initial concentration.

Solution

a) Using Eq. 1.42, $t_{1/2} = \frac{1}{k [A]_0} = \frac{1}{1.83 \times 10^{-3} \times 1.00 \times 10^{-2}}$
 $= 5.46 \times 10^4 \text{ s.}$

Time needed for the decrease in concentration of HI to one-eighth of its initial value is seven times the value calculated above as shown in Fig. 1.7; i.e. time needed is $3.82 \times 10^5 \text{ s.}$

Now we derive the integrated rate law for a reaction that is second order overall and is first order in each of the two reactants having different initial concentrations.

Case (ii): Integrated Rate Law for a Reaction that Follows Rate Law as per Eq. 1.35.

Let us start the reaction between A and B with different initial concentrations, $[A]_0$ and $[B]_0$. Let these two react to give products as per the rate law given in Eq.1.35



$$\text{Reaction rate} = k[A][B] \quad \dots(1.35)$$

For simplicity we have taken the reaction in which the stoichiometric coefficients of the reactants are same. But similar methods can be followed for reactions with different stoichiometric coefficients.

As per the reaction stoichiometry, A and B react in the ratio 1:1. After t seconds let the concentrations of A and B be $[A]_0 - x$ and $[B]_0 - x$.

Hence, the reaction rate = $\frac{-d[A]}{dt} = \frac{-d[B]}{dt} = k[A][B]$

i.e. $= \frac{-d\{[A]_0 - x\}}{dt} = k\{[A]_0 - x\}\{[B]_0 - x\}$

i.e. $= \frac{-d\{[A]_0\}}{dt} + \frac{dx}{dt} = k\{[A]_0 - x\}\{[B]_0 - x\}$

So $\frac{dx}{dx} = k\{[A]_0 - x\}\{[B]_0 - x\} \dots (1.43)$

$\frac{d[A]_0}{dt} = 0$ since $[A]_0$ is a constant.

In order to get the integrated form, we have to transform Eq. 1.43 as shown below:

$$\frac{dx}{\{[A]_0 - x\}\{[B]_0 - x\}} = kdt \dots (1.44)$$

Let us write the expression, $\frac{1}{\{[A]_0 - x\}\{[B]_0 - x\}}$ as follows:

$$\frac{1}{\{[A]_0 - x\}\{[B]_0 - x\}} = \frac{p}{\{[A]_0 - x\}} + \frac{q}{\{[B]_0 - x\}} \dots (1.45)$$

i.e., $\frac{1}{\{[A]_0 - x\}\{[B]_0 - x\}} = \frac{p\{[B]_0 - x\} + q\{[A]_0 - x\}}{\{[A]_0 - x\}\{[B]_0 - x\}}$

In other words, $p\{[B]_0 - x\} + q\{[A]_0 - x\} = 1 \dots (1.46)$

Put $x = [A]_0$, then $p\{[B]_0 - [A]_0\} = 1$

or $p = \frac{1}{[B]_0 - [A]_0} \dots (1.47)$

Hence, $p = -\frac{1}{[A]_0 - [B]_0} \dots (1.48)$

Put $x = [B]_0$ in Eq. 1.46

Hence $q\{[A]_0 - [B]_0\} = 1$

or
$$q = \frac{1}{[A]_0 - [B]_0} \quad \dots(1.49)$$

From Eqs. 1.45, 1.48 and 1.49

$$\frac{1}{\{[A]_0 - x\}\{[B]_0 - x\}} = - \frac{1}{\{[A]_0 - [B]_0\}\{[A]_0 - x\}} + \frac{1}{\{[A]_0 - [B]_0\}\{[B]_0 - x\}} \dots (1.56)$$

We have adopted the **partial fraction procedure** in the above steps.

Using Eq. 1.50 in Eq. 1.44

$$- \frac{dx}{\{[A]_0 - [B]_0\}\{[A]_0 - x\}} + \frac{dx}{\{[A]_0 - [B]_0\}\{[B]_0 - x\}} = k dt \quad \dots (1.57)$$

You can see that the two terms in the L.H.S. of Eq 1.51 contain either $\{[A]_0 - x\}$ or $\{[B]_0 - x\}$ in the denominator. The splitting of the expression in L.H.S of Eq. 1.44 through partial fraction method facilitates usage of Formula.

At time = 0, $x = 0$ (“x” denotes the change in concentration of A or B due to reaction)

At time = t , $x = t$

$$\int_0^{x_t} \frac{dx}{\{[A]_0 - [B]_0\}\{[B]_0 - x\}} - \int_0^{x_t} \frac{dx}{\{[A]_0 - [B]_0\}\{[A]_0 - x\}} = k \int_0^t dt \quad \dots (1.57)$$

$$\frac{1}{[A]_0 - [B]_0} \left[- \ln ([B]_0 - x) \right]_0^{x_t} - \frac{1}{[A]_0 - [B]_0} \left[- \ln ([A]_0 - x) \right]_0^{x_t} = kt$$

$$\frac{-1}{[A]_0 - [B]_0} \{ \ln ([B]_0 - xt) - \ln[B]_0 \} + \frac{1}{[A]_0 - [B]_0} \{ \ln ([A]_0 - xt) - \ln[A]_0 \} = kt$$

$$\frac{1}{[A]_0 - [B]_0} \ln \frac{([A]_0 - xt) [B]_0}{([B]_0 - xt) [A]_0} = kt \quad \dots(1.54)$$

$$\log \frac{([A]_0 - xt) [B]_0}{([B]_0 - xt) [A]_0} = \frac{k}{2.303} \{ [A]_0 - [B]_0 \} t \quad \dots(1.55)$$

$$\log \frac{([A]_0 - xt)}{([B]_0 - xt)} = \log \frac{[A]_0}{[B]_0} + k \frac{([A]_0 - [B]_0)}{2.303} t \quad \dots(1.56)$$


By plotting $\log \frac{[A]_0 - xt}{[B]_0 - xt}$ against t , a straight line is obtained. The slope of the

straight line is equal to $\frac{k([A]_0 - [B]_0)}{2.303}$

$$k = \frac{2.303 \times \text{slope}}{([A]_0 - [B]_0)} \quad \dots (1.57)$$

The name saponification is derived from the fact that soaps are produced by the hydrolysis of esters of long-chain fatty acids using alkalies.

Let us discuss the saponification of ester as an example for the calculation of second order rate constant. The hydrolysis of ester by an alkali is called saponification. For example, the saponification of methyl acetate can be represented by the following equation:



The saponification rate is studied as follows. A reaction mixture is prepared with known concentrations of alkali and ester. At regular intervals, certain volume of the solution is withdrawn and titrated against standard HCl. From the titre values, the concentrations of the unreacted alkali and the unreacted ester could be determined at various time intervals.

The rate constant can be determined using,

- i) Eq. 1.40 and 1.41, if the initial concentrations of ester and alkali are same;
 - ii) Eqs. 1.56 and 1.57, if their initial concentrations are different
- Now study the following example.

Example 8

The saponification of methyl acetate using sodium hydroxide was studied at 298 K. The initial concentrations of the alkali and ester in the reaction mixture were both 1.00×10^{-2} M. The reaction rate was followed by titration of a definite volume of the reaction mixture with standard HCl. The concentrations of unreacted alkali, $[A]_t$, at various time intervals are given below:

Time / s	240	550	720	1000	1550
$10^3[A]_t/\text{M}$	6.85	4.81	4.17	3.38	2.49

Calculate the second order rate constant.

Solution

Since the initial concentrations of the alkali and the ester are same, we can use

Eqs. 1.40 and 18.41 for solving this problem. We tabulate $\frac{1}{[A]_t}$ values

against various time intervals as follows:

$\frac{1}{[A]_t}$ M	146	208	240	296	402
t/s	240	550	720	1000	1550

As suggested by Eq. 1.40, $1/[A]_t$ against t plot is a straight line (Fig. 1.8) showing that the application of methyl acetic follows second order kinetics.

Using Eq. 1.41, $k = \text{Slope}$
 $= 0.194 \text{ M}^{-1} \text{ s}^{-1}$




Fig. 1.8: $1/[A]_t$ against t plot for the saponification of methyl acetate.

Having some experience with the differential and the integrated forms of rate laws for first order and second order reactions, we shall discuss zeroth order reactions in the next section. Also, we shall give some examples of third order reactions.

SAQ 4

State the units of the rate constants for zeroth order, first order and second order reactions. The rate of reaction is measured in $M s^{-1}$

.....
.....
.....

SAQ 5

The decomposition of HI is a second order reaction. At 700 K the rate constant for the reaction is $1.83 \times 10^{-3} M^{-1} s^{-1}$. If the initial concentration of HI is $1.00 \times 10^{-2} M$, calculate its concentration after $1.68 \times 10^5 s$.


.....
.....
.....

1.5 Summary

1.6 Terminal question

- 1) In the formation of ammonia,
$$N_2 (g) + 3H_2 (g) \rightarrow 2NH_3 (g)$$

the rate of consumption of hydrogen at a particular instant is $4.78 \times 10^{-4} M s^{-1}$. What is the rate of formation of ammonia?
- 2) The rate constant for the decomposition of N_2O_5 at 340 K is $5.20 \times 10^{-3} s^{-1}$. This reaction follows first order kinetics. If the initial rate of decomposition of N_2O_5 is $2.60 \times 10^{-4} M s^{-1}$, calculate the initial concentration N_2O_5 .
- 3) What is the time required for 87.5% decomposition of N_2O_5 at 340 K? Use data from the previous question.
- 4) What is the half-life period for the first order decomposition of azomethane at 600 K
if $k = 3.55 \times 10^{-4} s^{-1}$?



was studied at 298 K. The following results were obtained where

$$\text{rate} = \frac{\Delta[\text{S}_2\text{O}_8^{2-}]}{\Delta t}$$

$[\text{I}^-]_0$ and $[\text{S}_2\text{O}_8^{2-}]_0$ denote the initial concentrations of the two species

$10^2 \times [\text{I}^-]_0/\text{M}$	$10^2 \times [\text{S}_2\text{O}_8^{2-}]_0/\text{M}$	$10^5 \times \text{Initial rate}/\text{M s}^{-1}$
8.0	4.0	12.50
4.0	4.0	6.25
4.0	2.0	3.12

Determine the rate law.

1.7 Answer

SAS

1) Using Eq. 1.2 $\frac{d[\text{O}_2]}{dt} = \text{Rate}$
 $= 2.74 \times 10^{-4} \text{ M s}^{-1}$

2) $\frac{1}{2} \left(\frac{-d[\text{HI}]}{dt} \right) = \frac{d[\text{H}_2]}{dt}$

- 3) i) Spectrophotometry
 ii) pH measurement, conductance or titration
 iii) pressure measurement

4) Units of k :

zeroth order	:	M s^{-1}
first order	:	s^{-1}
second order	:	$\text{M}^{-1} \text{s}^{-1}$

5) Substituting in Eq. 1.40

$$\frac{1}{[\text{A}]_t} = \left[\frac{1}{[1.00 \times 10^{-2}]} + 1.83 \times 10^{-3} \times 1.68 \times 10^5 \right] \text{ M}^{-1}$$

$$[\text{A}]_t = 2.45 \times 10^{-3} \text{ M}$$

TMA

$$1) \quad \frac{1}{3} \frac{-d[\text{H}_2]}{dt} = \frac{1}{2} \frac{d[\text{NH}_3]}{dt}$$

Hence,
$$\frac{d[\text{NH}_3]}{dt} = \frac{2}{3} \left(\frac{-d[\text{H}_2]}{dt} \right) = \frac{2}{3} \times 4.78 \times 10^{-4} \text{ M s}^{-1}$$

$$= 3.19 \times 10^{-4} \text{ M s}^{-1}$$

$$2) \quad \frac{-d[\text{N}_2\text{O}_5]}{dt} = k [\text{N}_2\text{O}_5]$$

Hence,
$$[\text{N}_2\text{O}_5] = \frac{1}{K} \frac{-d[\text{N}_2\text{O}_5]}{dt} = \frac{2.60 \times 10^{-4} \text{ M s}^{-1}}{5.20 \times 10^{-3} \text{ s}^{-1}}$$

$$3) \quad [\text{A}]_t = 12.5\% [\text{A}]_0 = [\text{A}]_0/8$$

Using Eq. 1.20
$$t = \frac{2.303}{k} \log \frac{[\text{A}]_0}{[\text{A}]_0/8}$$

$$= \left(\frac{2.303}{5.20 \times 10^{-3} \log 8} \right) \text{ s} = 400 \text{ s}$$

$$4) \quad \text{Using Eq. 1.30 } t_{1/2} = \frac{0.693}{k} = \frac{0.693}{3.55 \times 10^{-4}} \text{ s} = 1950 \text{ s}$$

5) Let us write the rate law as,

$$v = k [\text{I}^-]^m [\text{S}_2\text{O}_8^{2-}]^n$$

As in Example 11,
$$m = \frac{\log 12.50 \times 10^{-6} / 6.35 \times 10^{-6}}{\log 8.0 \times 10^{-2} / 4.0 \times 10^{-2}} = 1$$

$$n = \frac{\log 6.25 \times 10^{-6} / 3.12 \times 10^{-6}}{\log 4.0 \times 10^{-2} / 2.0 \times 10^{-2}} = 1$$

The rate law is,

$$\text{rate} = k [\text{I}^-] [\text{S}_2\text{O}_8^{2-}]$$

UNIT 2 HIGHER ORDER OF REACTION

- 2.1 Introduction
- Objective
- 2.2 Zeroth Order
- 2.3 Third Order Reactions
- 2.4 Pseudo First Order Reactions
- 2.5 Determining the Order of Reaction
- 2.6 Fast Reactions
- 2.7 Summary
- 2.8 Terminal question
- 2.9 Answer

2.1 Introduction

In this unit we will study zeroth and third order equations. These equations will be used for calculating the rate constants of the reactions. We shall then explain the methods of determining the order of reaction.

Finally, the methods of studying fast reactions will be dealt with.

Objective

- derive integrated rate laws for zeroth and third order reactions and use them for calculating rate constants,
- state the methods for determining the order of reaction

2.2 ZEROth ORDER

The rate law for a zeroth order reaction is of the following form,

Zeroth order reaction is generally a heterogeneous reaction.


$$-\frac{d[A]}{dt} = k[A]^0 = k \quad \dots (1.58)$$

Since $[A]^0 = 1$


In a zeroth reaction, the reaction rate is independent of the concentrations of the reactant.

Some examples of zeroth order reactions are given below


- i) Decomposition of ammonia on a hot platinum surface,



ii) Decomposition of nitrous oxide on a hot platinum surface.



ii) Decomposition of hydrogen iodide on finely divided gold at 320 K.



Let us derive the integrated rate law for a zeroth order reaction.

Integrated Rate Law for Zeroth Order Reaction

Let us consider the zeroth order reaction:



Let the concentration of A at the start be $[\text{A}]_0$ and its concentration at the time t be $[\text{A}]_t$.

The integrated form of Eq.1.58 can be derived as follows:

$$\int_{[\text{A}]_0}^{[\text{A}]_t} -d[\text{A}] = \int_0^t k dt \quad \dots (1.59)$$

$$- \left\{ [\text{A}] \right\}_{[\text{A}]_0}^{[\text{A}]_t} = k(t - 0)$$

$$\begin{aligned} \text{i.e} \quad & -[\text{A}]_t + [\text{A}]_0 = kt \\ \text{or} \quad & [\text{A}]_0 - [\text{A}]_t = kt \\ & [\text{A}]_t = [\text{A}]_0 - kt \quad \dots (16.0) \end{aligned}$$

On plotting $[\text{A}]_t$ against t , a straight line is obtained for a zeroth order reaction. The slope is equal to $-k$

$$K = \text{Slope} \quad \dots (1.61)$$

Example 1

The decomposition of hydrogen iodide on gold at 323 K is zeroth order reaction and the rate constant is $1.20 \times 10^{-4} \text{ Ms}^{-1}$

- a) If the initial concentration of hydrogen iodide is 0.500 M, calculate its concentration after $3.00 \times 10^3 \text{ s}$.
- b) How long will it take for all of the hydrogen iodide to decompose?

Solution

(a) Using Eq. 1.60, $[A]_t = [A]_0 - kt$
 $= (0.500 - (1.20 \times 10^{-4} \times 3.00 \times 10^3)) \text{ M}$
 $= 0.140 \text{ M}$

(b) If hydrogen iodide completely decomposes, then $[A]_t = 0$
or $t = \frac{[A]_0}{k} = \frac{0.500 \text{ M}}{120 \times 10^{-4} \text{ M s}^{-1}}$
 $= 4.17 \times 10^3 \text{ s}$

Hence the reaction will be complete after $4.17 \times 10^3 \text{ s}$.

Half-Life of a Zeroth Order Reaction

As discussed for first and second order reactions,

$$[A]_t = [A]_0/2 \text{ when } t = t^{1/2},$$

Hence, Eq. 1.60 becomes,

or $[A]_0/2 - [A]_0 = kt^{1/2}$
 $kt^{1/2} = [A]_0/2$
 $t^{1/2} = \frac{[A]_0}{2k} \quad \dots (1.62)$

This means that the half-life of a zeroth order reaction is directly proportional to the initial concentration of the reactant.

Example 2

Calculate the half-life for the decomposition of hydrogen iodide on gold at 323 K. Use the data from Example 9.

Solution

Using the data from Example 9 in Eq.18.62,

$$t_{1/2} = \frac{[A]_0}{2k} = \frac{0.500 \text{ M}}{2 \times 1.20 \times 10^{-4} \text{ M}^{\text{s}^{-1}}} = 2.08 \times 10^3$$

From Examples 9 and 10, you can understand that the zeroth order reaction is complete in two half-lives as shown in Fig. 1.9





Fig. 1.9: Completion of a zeroth order reaction; t_1 is the time taken for both the conversions, $[A]_0 \rightarrow 50\% [A]_0 \rightarrow 0\% [A]_0$

2.3 Third Order Reactions

There are a few third order reactions. Two examples are given below:



The methods of arriving at differential and integrated rate laws for third order reactions are similar to those of first and second order reactions. We are not going to discuss the same.

SAQ 1

Using the data in Example 9, calculate the time required for the decomposition of HI on told at 323 K to proceed to 75%.


.....

.....

.....

2.4 PSEUDO FIRST ORDER REACTIONS

One of the ways of simplifying the study of reactions involving more than one reaction is to study under pseudo first order conditions. For example, considered the reaction:



$$\text{Reactions rate} = k [\text{CH}_3\text{Br}] [\text{OH}^-] \quad \dots(1.64)$$

If the concentrations of CH_3Br and OH^- are comparable, then the reaction is second order overall as indicated by Eq. 1.46. Suppose that the concentration of OH^- is much larger (say, 10 times or more) than the concentration of CH_3Br . In such cases the concentration OH^- does not change much during the reaction and can be considered constant. Hence, the reaction rate depends on the concentration of CH_3Br only.

$$\text{Reaction rate} = k' [\text{CH}_3\text{Br}]$$

Where $k' = k[\text{OH}^-]$; k' is the pseudo first order rate constant. The reaction can be treated as first order for calculation purposes. Hence, the integrated rate law is similar to Eq 1.21


$$\log \frac{[\text{CH}_3\text{Br}]_0}{[\text{CH}_3\text{Br}]_t} = \frac{k' t}{2.303}$$

In general, the reactions like the above which are effectively first order due to large excess of one of the reactions are called pseudo first order reaction.

Let us discuss the following two reactions which are studied under pseudo first order condition.

i) Acid Hydrolysis of Ester

The hydrolysis of ethyl acetate in presence of a mineral acid (say, HCl) can be represented by the following equation:



The reaction rate depends on [ester], [water] and $[\text{H}_3\text{O}^+]$. Here H_3O^+ ion is a catalyst. Since the concentration of the catalyst does not change during the reaction, and water is present in large amount, the reaction becomes pseudo first order in ester.

$$\text{Rate} = k' [\text{ester}]$$

where k' includes concentration and H_3O^+ . If the reaction is carried out a solvent other than water, the first order dependence on [water] also could be seen.

The pseudo first order rate constant is determined by titrating a definite volume of the reaction mixture containing ester and to HCl with standard alkali. Let V_0 , V_t and V_∞ be the volumes of standard alkali at the start, after a time t and after the completion of the reaction.

Infinite reading (V_∞) is usually taken after heating the reaction mixture for a few minutes or after keeping the reaction mixture at the experimental temperature for a long time.

V_∞ = Volume of alkali equivalent to i) acetic acid liberated after the completion of the reaction and ii) HCl present.

V_t = Volume of alkali equivalent to i) acetic acid and produced at the time t and ii) HCl present.

V_0 = Volume of alkali equivalent to HCl only.

Since the concentration of HCl is constant throughout the experiments,

$[\text{A}]_0$ (i.e, Initial concentration of ester) $\propto (V_\infty - V_0)$

and $[\text{A}]_t$ (i.e Concentration of ester remaining unreacted at t) $\propto (V_\infty - V_t)$

We can calculate pseudo first order rate constant for the acid hydrolysis of ethyl acetate by using the following modified form of Eq. 18.21.


$$\frac{k' t}{2.303} = \log \frac{[\text{A}]_0}{[\text{A}]_t} = \log \frac{V_\infty - V_0}{V_\infty - V_t}$$

... (1.65)

where k' is the pseudo first order rate constant.

ii) Inversion of Sucrose

The hydrolysis of sucrose to form glucose and fructose in presence of mineral acid is similar to the acid hydrolysis of ester as far as the reaction kinetics is concerned.



Sucrose turns the plane-polarised light to the right, (i.e., it is dextro rotatory). Glucose also turns the plane-polarised light to the right, while fructose turns it to the left (i.e., it is laevo rotatory). On completion of the reaction, the reaction

mixture is laevo rotatory, since the angle of rotation is more for fructose than for glucose. To start with, the reaction mixture is dextrorotatory due to Sucrose. Thus, the completion of reaction (infinite reading) is marked by the change in the sign of rotation. Due to this reason, the reaction is called inversion of sucrose.

If r_0 , r_t and r_∞ are, the angles of rotation at the beginning, after time t ; and after completion of the reaction, then the pseudo first order rate constant (k') for the inversion of sucrose is given by:

$$k' = \frac{2.303}{t} \log \frac{r_0 - r_\infty}{r_t - r_\infty} \quad (1.66)$$

Eq. 18.66 is a modified form of Eq. 18.21, where $[A]_0$ is proportional to $(r_0 - r_\infty)$ and $[A]_t$ and is proportional to $(r_t - r_\infty)$.

SAQ 2

Give the details of the graphical method of obtaining the pseudo first order rate constant for

- i) acid hydrolysis of ethyl acetate
- ii) inversion of sucrose.


.....

2.5 DETERMINING THE ORDER OF REACTION

In order to write the rate law, we must know the order of reaction with respect to each reactant. In this section, we discuss some methods for determining the order of reaction.

1) Method of Initial Rates

The instantaneous rate of reaction extrapolated to the instant when the reagents were just mixed is called the initial rate of the reaction. Let us consider the reaction



Let the rate of reaction be represented as,

$$v = k [A]^m [B]^n$$

where the reaction is m^{th} order in A and n^{th} order in B. The rate constant for the reaction is k . We have to obtain the initial rates from atleast two experiments in which the initial concentrations of A (a_1 and a_2) are different while the initial concentration of B (b_1) is constant

$$\text{Rate in Experiment I} = ka_1^m b_1^n$$

$$\text{Rate in Experiment II} \quad v_2 = ka_2^m b_1^n$$

From the ratio $\frac{v_1}{v_2}$, we can calculate m , since a_1 and a_2 are known.

$$\frac{\text{Rate in Experiment I}}{\text{Rate in Experiment II}} = \frac{v_1}{v_2} = \frac{ka_1^m b_1^n}{ka_2^m b_1^n} = \left[\frac{a_1}{a_2} \right]^m \quad \dots (1.67)$$

Extrapolation is the process of extending a curve upto a desired x or y coordinate to obtain the corresponding y or x value.

Initial reaction rate, could be graphically arrived at by plotting the concentration of a reactant against time. The tangent to the concentration curve is drawn at the very start of the reaction and its slope is calculated. The negative of the slope value is the initial rate.

Taking logarithms we can write,

$$\log \frac{v_1}{v_2} = m \log \frac{a_1}{a_2}$$

Similarly, the rate is measured for one more experiment in which the initial concentration of A is a_2 and the initial concentration of B is b_2 .

So rate in Experiment III $v_3 = k a_2^m b_2^n$

$$\frac{\text{Rate in Experiment I}}{\text{Rate in Experiment II}} = \frac{v_1}{v_2} = \frac{ka_2^m b_1^n}{ka_2^m b_2^n} = \left[\frac{b_1}{b_2} \right]^n \quad \dots(1.69)$$

$$\log \frac{v_1}{v_2} = n \log \frac{b_1}{b_2}$$

Since v_1 , v_2 , b_1 and b_2 are known, n can be calculated. The overall reaction order = $m + n$. You can understand this method from the following example.

Example 3

For the reaction $\text{Cl}_2 (\text{g}) + 2 \text{NO} (\text{g}) \longrightarrow 2\text{NOCl} (\text{g})$ the initial concentration $[\text{Cl}_2]_0$ and $[\text{NO}]_0$ are given along with initial rates.

$$\text{Rate} = \frac{-d[\text{Cl}_2]}{dt}$$

$[\text{Cl}_2]_0/\text{M}$	$[\text{NO}]_0/\text{M}$	Initial rate/ Ms^{-1}
0.10	0.10	3.0×10^{-3}
0.20	0.10	6.0×10^{-3}
0.20	0.20	2.4×10^{-2}

Calculate (i) order of the reaction with respect to each of the reactants and the overall order: (ii) what is the rate law? (iii) calculate the rate constant.

Solution

i) We can write the rate law as, $k [\text{Cl}_2]^m [\text{NO}]^n$.

Similar to Esq. 18.68 and 18.70, we can write the logarithm ratios and, calculate m and n as follows:

$$v_1 = k (0.10)^m (0.10)^n = 3.0 \times 10^{-3} \text{ M s}^{-1}$$

$$v_2 = k (0.20)^m (0.10)^n = 6.0 \times 10^{-3} \text{ M s}^{-1}$$

$$v_3 = k (0.20)^m (0.20)^n = 2.4 \times 10^{-2} \text{ M s}^{-1}$$

$$\text{Using Eq. 1.68, } m = \frac{\log \frac{3.0 \times 10^{-3}}{6.0 \times 10^{-3}}}{\log \frac{0.10}{0.20}} = \frac{\log \frac{6.0}{3.0}}{\log \frac{0.20}{0.10}} = 1$$

$$\text{Using Eq. 1.70, } n = \frac{\log \frac{6.0 \times 10^{-3}}{2.4 \times 10^{-2}}}{\log \frac{0.10}{0.20}} = \frac{\log \frac{2.4}{6.0}}{\log \frac{0.20}{0.10}} = 1$$

Hence, the reaction order in NO and first order in Cl_2 . The overall order is $2 + 1 = 3$.

ii) The rate law is given below:

$$\text{Rate} = k [\text{Cl}_2] [\text{NO}]^2$$

$[\text{A}]_0/10^{-2} \text{ M}$	5.00	4.00
$t_{1/2} / \text{s}$	240	300

Calculate the order of reaction.

.....
.....
.....

2.6 FAST REACTIONS

Many reactions are so fast that ordinary experimental methods mentioned in Sec. 18.3 are inadequate to make measurement of reaction rates or the rate constants. Such reactions are called fast reactions and the half-life periods of fast reactions are less than 10^{-2} s. Some of the special techniques used for measuring the constants of fast reactions are:

- flash photolysis
- flow methods, and
- relaxation methods.

Of the above three, we shall discuss the first method in CHE 411. Now we consider the principles of the other two methods.

Flow Method

Two techniques are available under flow method. In continuous flow method (Fig. 1.10) the reacting solutions or gases are taken in separate containers (A and B) and are allowed to flow through the mixing chamber (C) into an observation tube (D). At various points along the observation tube, the composition of the mixture is determined by some physical methods.

iii) The rate constant can be calculated using any one of the three rates given above.


$$v_1 = 3.0 \times 10^{-3} \text{ M s}^{-1} = k (0.10 \text{ M}) (0.10 \text{ M})^2$$

$$k = \frac{3.0 \times 10^{-3}}{(0.10)^3} \text{ M}^{-2} \text{ s}^{-1} = 3.0 \text{ M}^{-2} \text{ s}^{-1}$$

Care must be taken in applying the method of initial rates. For complex reactions like the formation of HBr (discussed in the next section) the product also affects the rate. The method of initial rates is applicable to simple reactions only.

Clock Reactions


In the case of some reactions, the time taken for the colour change of the reaction mixture can be used for measuring the initial rate. Such self indicating reactions are known as clock reactions. For example, the kinetics of the reaction,



Potassium
perdisulphate

can be monitored as a clock reaction. A reaction mixture is prepared using potassium perdisulphate and potassium iodide in a higher concentration and sodium thiosulphate in much lower concentration. A group of (starch is also present in the reaction mixture. The reaction mixture colourless in the beginning and it turns blue after some time.

The time, Δt , between the mixing of the reactants and the appearance of blue colour noted. The blue colour develops due to the liberation of free iodine, after sodium thiosulphate (present in less concentration) is consumed completely as per the reaction:



Sodium
tetrathionate

The order of the reaction with respect to KI (m) and the order of the reaction with respect to $\text{K}_2\text{S}_2\text{O}_8$ (n) can be calculated by using the following formula:

$$\log (1/\Delta t) = m \log [\text{KI}] + n \log [\text{K}_2\text{S}_2\text{O}_8] + \text{constant}$$

A plot of $\log (1/\Delta t)$ against $\log [\text{KI}]$ is made using Δt values obtained by varying $[\text{KI}]$ and keeping $[\text{K}_2\text{S}_2\text{O}_8]$ constant. The slope of the straight line gives m. Similarly n is obtained from the slope of the straight line got by plotting $\log (1/\Delta t)$ against $[\text{K}_2\text{S}_2\text{O}_8]$. For the second plot, Δt is obtained by varying $[\text{K}_2\text{S}_2\text{O}_8]$ and keeping $[\text{KI}]$ constant.

Experimentally it has been found that $m = 1$ and $n = 1$

$$\text{Hence, } -d[\text{K}_2\text{S}_2\text{O}_8] = k [\text{KI}] [\text{K}_2\text{S}_2\text{O}_8]$$

For clock reactions, two more examples are given below:

- 1) Acid catalysed iodination of acetone
- 2) Saponification of ester (using phenolphthalein indicator).

Some methods of studying fast reactions will be discussed in Sec. 1.72.

2) Trial and Error Method

We can determine the order of reaction

- i) by substitution for experimental data into Eqs. 1.21, 1.40 and 1.60
- ii) by graphical method using plots such as $\log [A]$ against t , $1/[A]$ against t and $[A]$ against t . The order of the reaction is one, two or zero depending on
 - i) which of the equations gives rise to a constant value for k or
 - ii) which of the plots gives a straight line.

3) Half-Life Method

The half-lives are determined using different initial concentrations of the reactant. If the half-life is independent of initial concentration, the reaction is first order. If the half-life is inversely proportional to the power of initial concentration, the reaction is second order. If the half-life is directly proportional to the first power of initial concentration, the reaction is zeroth order.

In general, half-life period (t) is proportional to $[A]_0^{1-n}$ where $[A]_0$ is the initial concentration of the reactant and n is the order of the reaction.

If the half-life periods are t_1 and t_2 corresponding to the initial concentrations

$[A]_1$ and $[A]_2$ of a reactant, then

$$\frac{t_2}{t_1} = \left(\frac{[A]_2}{[A]_1} \right)^{1-n} = \left(\frac{[A]_1}{[A]_2} \right)^{n-1}$$
$$n-1 = \frac{\log t_2/t_1}{\log [A]_1/[A]_2}$$
$$n = 1 + \frac{\log t_2/t_1}{\log [A]_1/[A]_2} \quad \dots(1.71)$$

Example 4

For the decomposition of acetaldehyde in gas-phase at 791 K, the half-life periods are 328 s and 572 s corresponding to the initial concentration 9.72×10^{-3} M and 4.56×10^{-3} M.

Solution

Using Eq. 1.71


$$n = 1 + \frac{\log 572/328}{\log 9.72 \times 10^{-3}/4.56 \times 10^{-3}} = 1 + \frac{0.2415}{0.3287} = 1.735$$

The order of the reaction is 1.735. Note that the order of the reaction is fractional.

4. Isolation Method

In the case of reaction having more than one reactant, the rate law can be simplified if the concentrations of all reactants except one are taken in excess. The reaction rate then depends on the reactant present in lesser quantity. The order of the reaction is determined by one of the methods given above. It is equal to the order in the reactant present in lesser quantity. This procedure is repeated in turn with each of the reactants being in lesser amount while others are in excess. The procedure is called Van't Hoff's isolation method.

For example, consider the reaction,



For which the rate law is,

$$\text{Rate} = k [A]^m [B]^n \quad \dots (1.72)$$

In the first set of experiments B is in excess as compared to A, such that the rate depends on A only.

$$\text{Hence, the rate} = k_1 [A]^m \quad \dots (1.73)$$

$$\text{Where } k_1 = k (\text{Initial concentration of B})^n \quad \dots (1.74)$$

Since B is in excess as compared to A, the concentration of B almost remains a constant throughout the experiment. From the rate the measurements, m can be found out using graphical method.

In the second set of experiments, [A] is much large as compared to [B]. The rate measurements are made and using the rate law stated below, n is calculated.

$$\text{rate} = k_2[\text{B}]^n \quad \dots (1.75)$$

where $k_2 = k$ (Initial concentration of A)^m ... (176)

The overall order of the reaction is $m + n$.

SAQ 3

For the alkaline hydrolysis of ethyl nitrobenzoate, the half-life periods and the initial concentrations are given below:




Fig. 1.10 Apparatus for continuous flow method

Using the method, reactions which get completed within 10^{-3} s can be studied. But this method requires a large volume of the reactants. In order to study the reaction rates using small volumes of samples, a refined method is used which is known as stopped flow method. The reacting solutions are forced through jets into a mixing chamber where mixing occurs very rapidly (within 10^{-3} s.) The solution passes at once into the reaction vessel from the mixing chamber. The flow is then stopped and measurements are made using suitable physical methods. This method is widely used for the study of enzymes kinetics.

Relaxation Method

For studying reactions, which are complete within 10^{-5} s or even less, one of the problems encountered is that the time of mixing the reactants should be much smaller than the time taken for the completion of the reaction. To avoid this problem, relaxation methods have been developed. In a relaxation method, we disturb a reaction at equilibrium using an external influence such as sudden variation of pressure or, temperature (known as pressure jump or temperature jump). The system is said to be perturbed from its equilibrium position.

The perturbed system then returns to a new equilibrium position. Relaxation refers to the passage of a perturbed system to the new equilibrium. The concentration of the perturbed system is recorded at various time intervals, using suitable physical methods. From these measurements, it is possible to measure the rate constants.

2.7 SUMMARY

We derived the integrated forms of rate expressions for third order and zeroth order reactions. We defined and illustrated the pseudo first order reactions. We explained the methods of determination of order of reaction. Finally, we gave an outline of the methods] of studying fast reactions.

2.8 TERMINAL QUESTIONS

- 6) Explain two methods for the study of fast reactions.
- 7) The relaxation methods are suitable for measuring the rates of fast reactions. The experimental techniques (such as titremetry mentioned in Sec. 18.3) are not useful for this purpose. Explain the reason.

2.9 ANSWER

Self Assessment Questions

- 1) Using Eq. 1.60

$$[A]_t = \frac{25}{100} \times 0.500 \text{ M} = 0.125 \text{ M}$$

$$t = \frac{(0.500 - 0.125) \text{ s}}{1.20 \times 10^{-4}}$$
$$= 3.13 \times 10^3 \text{ s}$$

- 2) i) After obtaining V_0 , V_t and V_∞ as explained in sec. 1.6, $\log (V_\infty - V_t)$ is plotted against t .

$$k^I = -2.303 \times \text{slope}$$

where k^I is the pseudo first order rate constant for the acid hydrolysis of ester.

- ii) Using r_0 , r_t and $r =$ values, $\log (r_1 - r)$ is plotted against t .

$$k = -2.202 \times \text{slope}$$

where k' is the pseudo first order rate constant for the inversion of sucrose.

3) Using E.q 1.71 $n = 1 + \frac{\log 300/240}{\log 5.00 \times 10^{-2}/400 \times 10^{-2}}$

$$n = 1 + \frac{0.0969}{0.0969} = 2$$

Terminal Questions

1. See in Sec 1.11
2. Methods such as titrimetry need mixing of the reactants. In the case of many fast reactions, the reaction is faster than the mixing time of the reactants. Hence, methods such as titrimetry are not useful for measuring rates of fast reactions. Relaxation method avoids the problem of mixing the solutions, and hence, is useful for measuring the rates of fast reactions.

UNIT 3 SOME REACTION MECHANISMS

- 3.1 Introduction
 - Objectives
- 3.2 Some reaction mechanisms
- 3.3 Theory of Unimolecular Reactions
- 3.4 Theories of reaction rates
- 3.5 Summary
- 3.6 Terminal question
- 3.7 Answer
- Appendix

3.1 INTRODUCTION

Chemical kinetics is the study of rates and mechanisms of chemical reactions. The rate of a reaction depends on many factors such as the concentration of the reactants, temperature, catalysts, etc. Some of these factors were examined in unit 1.

We shall discuss the mechanism of simple reactions. We shall illustrate the significance of rate determining step in explaining the rate law. We shall study the effect of temperature on reaction rates. We shall state the theories of reaction rates.

Objectives

After studying this unit, you should be able to:

- define elementary reaction and molecularity,
- propose mechanism for simple reactions using the experimental rate law
- discuss the theory of unimolecular reactions,
- explain Arrhenius equation, collision theory and activated complex theory, and
- describe the methods of studying fast reactions.

3.2 SOME REACTION MECHANISMS

Many chemical reactions take place through a series of steps. Each step is known as an **elementary reaction**. A **reaction mechanism** is a series of elementary reactions proposed for explaining the rate law for the overall reaction. The elementary reactions are written as chemical equation. Such chemical equations give a possible explanation for the reaction path.

For an elementary reaction, the molecularity is the same as the order of reaction.
--

The rate law for each elementary reaction can be written using **molecularity**. The

molecularity is the number of reactant molecules or atoms in an elementary reaction. If there is only one reactant molecule (or atom) in an elementary reaction, the reaction is said to be unimolecular. An elementary reaction in which two molecules (or atoms) react is bimolecular reaction. Most reaction mechanisms consider mainly unimolecular and bimolecular reactions. The chance of termolecular reactions (where three species are to combine) occurring is much less, because the probability of three species colliding simultaneously is quite low. An example each for unimolecular and bimolecular reactions is given below.

Unimolecular Reaction



A unimolecular reaction has a first order rate law, hence the rate of decomposition of O_3 could be represented as follows;

$$\text{Rate} = k [\text{O}_3] \quad \dots(2.2)$$

Bimolecular Reaction



A bimolecular reaction has an overall second order rate law, being first order in each reactant. Hence, for the elementary reaction given in Eq. 18.79, rate can be expressed as follows:

$$\text{Rate} = k [\text{O}] [\text{O}_3] \quad \dots(2.4)$$




Some of the guidelines followed in suggesting reaction mechanisms are given below:

- 1) The elementary reactions when added must be equal to the overall balanced chemical equation for the reaction.

For example the overall reaction in the decomposition of O_3 in the upper atmosphere is,



This reaction could be taken as the result of the following two elementary reactions:



- 2) While writing such a mechanism, one possible support is proving the presence of **intermediates**. For example, in the mechanism suggested above, atomic oxygen is an intermediate. Such intermediates can be detected by physical or chemical methods. They are generally reactive species. Further an intermediate is produced and ultimately used up.
- 3) The mechanism must agree with the overall rate law determined; experimentally. In other words the rate laws for the elementary reactions must be combined in such a way that the overall rate law is explained. In order to accomplish this, we must be able to decide the **rate determining step**. Out of the elementary reactions suggested, the slowest one is called the rate determining step. The overall reaction rate cannot be faster than the slowest step in a mechanism. The rate determining step decides the rate of the overall reaction.

For example, in the mechanism suggested for the decomposition of ozone, Step ii (i.e., Eq 2.3) is possible the rate determining step. We shall explain this shortly.

- 4) The possibilities of both forward and reverse reactions occurring fast must also be considered! That is the possibility of a dynamic equilibrium must also be examined. This is one of the ways to
 - find a suitable relationship for expressing the concentration of an intermediate and
 - eliminate the term denoting the concentration of the intermediate from the rate expression for the overall reaction.
- 5) Kinetic information can only support a proposed mechanism; it should not be taken as a proof since a mechanism cannot be proved absolutely.

Only a few guidelines are given here for proposing a reaction mechanism. However, these are sufficient for studying the reaction mechanisms of simple reactions.

The studies on organic and inorganic reaction mechanisms have led to the growth of separate branches of chemistry. You are advised to take up the course on Organic Reaction Mechanisms, if you desire to study theories of organic reactions.

Now, we shall discuss the reaction mechanism involving:

- a fast equilibrium followed by a slow step.
- a slow step followed by a fast step, and

- a chain reaction.

We shall also state the following types with an example in each case without discussing the reaction mechanisms.

- Consecutive reactions,
- Opposing reactions; and
- Parallel reactions

Example 1

For the decomposition Of O_3 ,





the overall rate law is given below:

$$\text{Overall rate} = k \frac{[O_3]^2}{[O_2]} \quad \dots(2.6)$$

Suggest a possible mechanism to explain the observed rate law.

Solution

You can see that the overall rate contains the term for the concentration of oxygen which is a product. This indicates that the mechanism consists of more than one step. A possible mechanism is suggested through Eqs. 2.7 and 2.3



Note that the addition of Eq 2.3 and the forward reaction of Eq. 2.7 gives the overall balanced equation as per Eq. 18.81. In the mechanism suggested above, k_1 and k_{-1} are the rate constants for the forward and the reverse reactions as per Eq.18.83; k_2 is the rate constant for the reaction as per Eq. 2.3

As per the mechanism suggested above,

$$\text{Overall rate} = k_2 [\text{O}_3] [\text{O}] \quad \dots(2.4)$$

Now we shall eliminate [O] in Eq. 18.80. For this purpose, we shall assume the following

$$\text{Rate of formation of O} = \text{Rate of Consumption of O} \quad \dots(2.8)$$

$$\text{i.e.,} \quad k_1 [\text{O}_3] = k_{-1}[\text{O}_2][\text{O}] + k_2[\text{O}_3][\text{O}] \quad \dots(2.9)$$

This assumption is valid because [O] is extremely small at any given time. Its variation with time ($d[\text{O}]/dt$) is still small and it may be taken to be zero. This type of assumption is called steady state approximation and we shall discuss it in Sec. 2.3

The above equation is written in the basis of the proposed mechanism which implies that oxygen atom is formed in the forward reaction of step (i) and is consumed in the reverse reaction of step (i) and, in step (ii). The forward and reverse reaction as per step (i) are much faster than the reaction as per step (ii). In other words, $k_2 [\text{O}_3] [\text{O}]$ is negligibly smaller than $k_{-1}[\text{O}] [\text{O}]$.

Hence Eq. 2.9 becomes,

$$k_1 [\text{O}_3] = k_{-1}[\text{O}_2] [\text{O}] \quad \dots(2.10)$$

$$\text{or} \quad [\text{O}] = \frac{k_1 [\text{O}_3]}{k_{-1} [\text{O}_2]} \quad \dots(2.11)$$

Using Eq 2.11 in Eq 2.4

$$\begin{aligned} \text{Overall rate} &= k_2[\text{O}_3][\text{O}] \\ &= \frac{k_1 k_2}{k_{-1}} \frac{[\text{O}_3][\text{O}_3]}{[\text{O}_2]} \quad \dots(2.12) \end{aligned}$$

$$\text{i.e.,} \quad \text{Overall rate} = k \frac{[\text{O}_3]^2}{[\text{O}_2]} \quad \dots(2.13)$$


where k is the composite rate constant since it combines the rate constants, k_1 , k_2 and k_{-1} per the equation,

$$k = \frac{k_1 k_2}{k_{-1}} \quad \dots(2.14)$$

You can see that Eq. 2.13 is the same as Eq 2.6

Example 2

The rate expression for the reaction,



is given below:




$$\text{Overall rate} = k [\text{NO}_2]^2$$

Spectroscopically NO_3 radical has been detected. Suggest a mechanism in keeping with these facts.

Solution

While suggesting the mechanism, the following facts must be borne in mind:

- i) The stoichiometry for the overall reaction must be as per Eq 2.18
- ii) As per Eq 2.19, the rate does not depend on $[\text{CO}]$ but depends on $[\text{NO}_2]^2$. This means that the reaction consists of more than one step and CO does not take part rate determining step.
- iii) NO_3 is a probable intermediate in the reaction. Keeping in view the above features, the following mechanism is proposed:



The stoichiometry is as per Eq. 2.18. Since $[\text{CO}]$ does not appear in the rate expression step (i) is assumed to be slower than step (ii). It is almost assumed that the intermediate, NO_3 , is consumed at a faster rate than it is formed. In other words.

i.e., Overall rate = Rate of formation of NO_3

$$\left. \begin{array}{l} \text{But rate of formation of} \\ \text{NO}_3 \text{ (as per Eq. 2.20)} \end{array} \right\} = k [\text{NO}_2]^2$$

which is same as Eq. 2.19

Example 3

The rate for the reaction between hydrogen and bromine is quite complex as given by Eq 2.23




$$\frac{d [\text{HBr}]}{dt} = \frac{k [\text{H}_2] [\text{Br}_2]^{1/2}}{1 + \frac{k' [\text{HBr}]}{[\text{Br}_2]}} \quad \dots(2.23)$$

Suggest a mechanism which could explain in the above rate law.

Solution

To explain the rate law a reaction sequence is proposed as follows:



The rate of reaction can be represented by Eq. 18.102 noting that HBr is formed as per Eqs. 2.25 and 2.26 and consumed as per Eq. 2.27.

$$\frac{d [\text{HBr}]}{dt} = k_2 [\text{Br}][\text{H}_2] + k_3 [\text{H}] [\text{Br}_2] - k_4 [\text{H}] [\text{HBr}] \quad \dots(2.29)$$

The negative sign preceding $k_4 [\text{H}] [\text{HBr}]$ in Eq. 2.29 is due to the consumption of HBr as per Eq. 2.27. The reaction sequence given above is an example of a chain reaction. In a chain reaction, an intermediate reacts to produce another intermediate. For example, Br, a radical intermediate produced as per Eq. 2.24

reacts with H₂ to produce another radical intermediate, H. Again H reacts with Br₂ as per Eq. 2.26 to give Br and so on. Reaction represented by Eq. 2.24 is the **chain initiation** reaction, since it is the start of the chain reaction. Reactions such as Eqs 2.25-2.327 are known as **chain propagation** reactions. Although Eq. 2.27 represents a propagation reaction producing Br, it also results in the consumption of HBr. For the latter reason, Eq. 2.27 represents an **inhibiting reaction**. Eq. 2.28 is a **chain terminating reaction** since the intermediates combine to give a molecule.

Often an intermediate in a chain reaction is a free radical. A free radical is an atom or a fragment of a molecule and has an unpaired electron.

To simplify Eq. 2.29, we must express [H] and [Br] in terms of [Br₂], [H₂] and [HBr]. Such a step is required since the concentrations of the intermediates are not easy to measure whereas the concentrations of the reactants and products could be measured. Such a simplification is possible assuming that the net rates of formations of intermediates are equal to zero.

i.e.,
$$\frac{d[\text{Br}]}{dt} = 0 \quad \dots (2.30)$$

and
$$\frac{d[\text{H}]}{dt} = 0 \quad \dots(2.31)$$

As per the elementary reactions given above, Br is formed as per Eqs. 2.24, 2.26 and 2.27; Br is consumed as per Eqs. 2.25 and 2.28. Also two bromine atoms are as per Eq. 2.24 for every bromine molecule dissociated and two bromine atoms are consumed as per Eq. 2.28 for every bromine molecule formed. Using these ideas and q. 2.26, we can write,

$$\frac{d[\text{Br}]}{dt} = 0 = 2k_1 [\text{Br}_2] - k_2 [\text{Br}][\text{H}_2] + k_3 [\text{H}][\text{Br}_2] + k_4 [\text{H}] [\text{HBr}] - 2k_5[\text{Br}]^2 \quad \dots(2.32)$$

Similarly, H is formed as per Eq. 18.98 and consumed as per Eqs 2.26 and 2.27. Using these ideas and Eq.2.31 we can write,

$$\frac{d[\text{H}]}{dt} = 0 = k_2 [\text{Br}][\text{H}_2] - k_3 [\text{H}][\text{Br}_2] - k_4 [\text{H}] [\text{HBr}] \quad \dots(2.33)$$

By adding Eqs. 2.32 and 2.33 we get,

$$2k_1 [\text{Br}_2] = 2k_5 [\text{Br}]^2$$

or
$$[\text{Br}] = \left(\frac{k_1}{k_5} \right)^{1/2} [\text{Br}_2]^{1/2} \quad \dots (2.34)$$

From Eq. 2.33 we can write,

$$[\text{H}] = \frac{k_2 [\text{H}_2] [\text{Br}]}{k_3 [\text{Br}_2] + k_4 [\text{HBr}]} \quad \dots(2.35)$$

Using Eqs. 2.34 and 2.35, we get

$$[\text{H}] = \frac{k_2 \left(\frac{k_1}{k_5}\right)^{1/2} [\text{H}_2] [\text{Br}_2]^{1/2}}{k_3 [\text{Br}_2] + k_4 [\text{HBr}]} \quad \dots(2.36)$$

Rearranging Eq. 2.29, we get,

$$\frac{d[\text{HBr}]}{dt} = k_2 [\text{H}_2][\text{Br}] + [\text{H}] (k_3[\text{Br}_2] - k_4 [\text{HBr}]) \quad \dots(2.37)$$

Using Eqs. 2.34, 2.36 and 2.37, we get

$$\begin{aligned} \frac{d[\text{HBr}]}{dt} &= k_2 \left(\frac{k_1}{k_5}\right)^{1/2} [\text{Br}_2]^{1/2} [\text{H}_2] + \frac{(k_3 [\text{Br}_2] - k_4 [\text{HBr}]) (k_2 (k_1/k_5)^{1/2} [\text{H}_2] [\text{Br}_2]^{1/2})}{k_3 [\text{Br}_2] + k_4 [\text{HBr}]} \\ &= \frac{2k_2 k_3 (k_1/k_5)^{1/2} [\text{Br}_2]^{1/2} [\text{H}_2]}{k_3 [\text{Br}_2] + k_4 [\text{HBr}]} \quad \dots (2.38) \end{aligned}$$

Dividing the numerator and denominator of R.H.S by $k_3 [\text{Br}_2]$,

$$\frac{d[\text{HBr}]}{dt} = \frac{2k_2 (k_1/k_5)^{1/2} [\text{H}_2] [\text{Br}_2]^{1/2}}{\frac{1 + k_4 [\text{HBr}]}{k_3 [\text{Br}_2]}} \quad \dots (2.39)$$

By comparison you can see that Eqs. 2.23 and 2.39 are same, where

$$k = 2k_2 \left(\frac{k_1}{k_5}\right)^{1/2} \text{ and } k' = \frac{k_4}{k_3}$$

The presence of the term $[\text{HBr}]/[\text{Br}_2]$ in the denominator of Eq. 2.23 or Eq 2.39, has the following significance.


- HBr formed inhibits the rate of the reaction
- At high $[\text{Br}_2]$, the inhibition is less

Both these predictions have been verified experimentally.

In the H_2 - Br_2 reaction, inhibition takes place due to the product formed. Sometimes, impurities are deliberately added to some materials to inhibit undesirable reactions. For example, food products generally get spoiled due to chain reactions involving oxidation. To preserve the food products, it is customary to use the preservatives which inhibit oxidation by removing chain-propagating radicals. Such "antioxidants" are also added to plastics and rubber to prevent their degradation.

Free-Radical Reactions

In H_2 - Br_2 reaction, H and Br atoms have unpaired electrons and these are free-radicals. In 1929, Paneth and Hofeditz reported the formation of polyatomic free radicals (CH_3 radicals) by the thermal decomposition of lead tetramethyl. It was found that lead was deposited as a mirror, in the hot portion of a tube through which hydrogen gas carrying lead tetramethyl vapour was passed.



The free-radicals such as CH_3 and C_2H_5 are detected by direct experimental methods or from the products they give. Experimentally obtained rate law is given by the equation.

$$\text{rate} = k [C_2H_6] \quad \dots (2.41)$$

where k is the overall rate constant: k is the complex combination of the rate constants of the individual elementary reactions.

Consecutive Reactions

We have postulated the existence of intermediates. In many cases, the intermediate in one step is the reactant in the next. Such reactions are called consecutive reactions. The rates of consecutive reactions could be stated in terms

of the concentrations of the reactants taken initially and the products formed in each stage.

Example

Acid hydrolysis of diethyl adipate.

The radical intermediates can be removed by using substances like NO. Since NO molecule has an unpaired electron, it combines with a radical intermediate which also has an unpaired electron. This could result in chain termination. Here NO molecule is called the radical scavenger and it is said to quench the chain reaction. To prove the chain mechanism, such radical scavengers are used.


Just as we can terminate a chain reaction using radical scavengers, we can start a chain reaction using free radical sensitizers such as Pb(CH₃)₂ or Hg(CH₃)₂. To increase the decomposition rate of an organic compound, Pb(CH₃)₂ or Hg(CH₃)₂ is added. These substances decompose, and introduce CH₃ radicals into the system. This starts the decomposition of the organic compounds through a chain reaction. Pb(CH₃)₄ and Hg(CH₃)₂ are said to sensitize the decomposition of organic compounds.

Opposing Reactions

In opposing reactions, rates of forward and reverse reactions are both appreciable. While proposing a mechanism both the reaction rates must be considered.

Example

Formation of butyrolactone from γ -hydroxybutyric acid.




Parallel Reactions

If a reactant can undergo more than one reaction, the resulting reactions are called parallel reactions. The rates of a set of parallel reactions can be measured concentrations of the products formed in each case.

Example




Nitration of phenol yielding *o*-nitrophenol and *p*-nitrophenol,



In the next section we shall explain the theory of unimolecular reaction rates.

SAQ 1


For the following elementary reactions, write the rate laws:



.....
.....
.....

SAQ 2



For the reaction,



the rate law is given below:

$$\text{Rate} = k[\text{NO}]^2 [\text{O}_2]$$

Is the following mechanism consistent with the rate law?



Hint: Consider NO₃ as an intermediate.

.....
.....
.....

3.3 THEORY OF UNIMOLECULAR REACTIONS

A unimolecular reaction is an elementary reaction in which only one molecule or a radical reacts. The unimolecular reactions follow first order kinetics. A number of gas phase reactions follow first order kinetics. These reactions are assumed to proceed through unimolecular rate-determining step. But how does the reactant molecule attain the activation energy? The activation energy is the minimum energy needed do the reactant molecules to react and yield products. If the molecules obtain their activation energy through collisions, it is difficult to explain first order kinetics. A collision process needs atleast two molecules and hence, second order kinetics could be expected but not first order kinetics. In 1992, Lindemann and Hinshelwood proposed a mechanism which could explain the unimolecular reactions in which molecules attain their activation energy through collision.


Let two molecules of the reactant gas (X) collide yielding an activate molecule (X^*) and a normal molecule (X). Such a collision is called an activating collision.



$$\text{Rate of activation of } X = k_a [X] [X] \quad \dots (2.45)$$

The activated molecule, X^* , can undergo either of the following reactions:

- i) X^* can undergo collision with another molecule X and lose its excess energy. Such a collision is called a deactivating collision



$$\text{Rate of decay of } X^* = k_a' [X^*] [X]$$

- ii) Alternatively, of X^* can decay to yield the product, Y .



$$\text{Rate of decay of } X^* = \text{Rate of product formation} = \frac{d[Y]}{dt} = k_b [X^*] \quad \dots(2.47)$$

Note that decay of X^* is a unimolecular reaction.

If the decay of X^* yielding products is the rate-determining step, then the overall rate of the reaction is given by the following expression.

$$\text{Rate} = k_b [X^*] \quad \dots(2.48)$$

In order to express the concentration of X^* , an active species, in terms of concentrations of reactants (or products) in the ground state, **steady-state approximation** is used. According to this procedure, it is assumed that a steady-state is reached after a reaction starts such that the concentration of the activated species is more or less a constant and does not change with time.

$$\text{i.e.} \quad \frac{d[X^*]}{dt} = 0 \quad \dots(2.49)$$

According to steady-state approximation, the concentrations of all reactive intermediates are constant and small during the major part of the reaction.

This means that the activated species, X^* , is consumed as soon as it is formed. Since X^* is formed as per Eq. 18.15 and is consumed as per Eqs. 2.44 and 2.46,

$$\frac{d[X^*]}{dt} = 0 = k_a [X]^2 - k_4 [X] [X^*] - k_b [X^*] \quad \dots(2.50)$$

$$\text{i.e.,} \quad [X^*] (k_4 [X] + k_b) = k_a [X]^2$$

$$\text{or} \quad [X^*] = \frac{k_a [X]^2}{(k_4 [X] + k_b)} \quad \dots(2.51)$$

Using Eqs. 2.48 and 2.51

$$\text{rate} = \frac{k_a k_b [X]^2}{k_4 [X] + k_b} \quad \dots(2.52)$$

From $p = cRT$ or $p \propto c$. Hence, when the pressure of a gas is high, its concentration is high. When the concentration is high, there will be large number of collisions.

$$(k_4 [X] + k_b)$$

At High Pressures

At high pressures, the number of collisions is large and the probability of deactivating collisions occurring; is high that is the rate of deactivation is larger than the rate of product formation (through decay); the unimolecular decay of X^* is the rate-determining step at high pressures; i.e.

Eqs. Such as 2.54 or 2.37 are instances of limiting conditions. Using such approximation procedure, it is possible to simplify a complicated equation as Eq. 2.52.

$$k'_a [X^*] [X] \gg k_b [X^*]$$

or $k'_a [X] \gg k_b$... (2.53)

In other words, $k'_a [X] + k_b \simeq k'_a [X]$... (2.54)

Using this in Eq. 2.52, we get

$$\text{rate} = \frac{k_a k_b [X]^2}{k'_a [X]} = \frac{k_a k_b [X]}{k'_a} \quad \dots (2.55)$$

In other words, the rate is first order at high pressures.

At Low Pressures

At low pressures, the number of collisions decreases. This means that the activated molecule yields the product as soon it is formed and there is not much time left for deactivating collision to occur. In other words, the bimolecular formation of X^* is the rate-determining step. Further, the rate of deactivating collisions is much small as compared to the rate of product formation.

$$k_b [X^*] \gg k'_a [X^*] [X]$$

or $k_b \gg k'_a [X]$... (2.56)

or $k'_a [X] + k_b \simeq k_b$... (2.57)

Using this in Eqs. 18125,

$$\text{rate} = \frac{k_a k_b [X]^2}{k_b} = k_a [X]^2 \quad \dots (2.58)$$

Hence the reaction follows second order kinetics at low pressures,

The variation of the order of reaction with pressure as predicted by Lindemann - Hinshelwood theory could be observed in reactions such as the decomposition of N_2O_5 at high pressures. Using this theory we could explain the unimolecular decomposition of N_2O_5 at high pressures.

3.4 THEORIES OF REACTION RATES

The rates of many reactions increase with the rise in temperature. Arrhenius proposed the following empirical relationship between the rate constant, k , and temperature, T .

$$\ln k = \ln A - E_a/RT \quad \dots(2.59)$$

or
$$\log k = \log A - E_a/2.303RT \quad \dots(2.60)$$

where A is called the Arrhenius factor or frequency factor or pre-exponential factor and E_a is the activation energy. Activation energy is the threshold energy that the reactant molecules must have in order to react. If $\log k$ is plotted against $\frac{1}{T}$, a straight line (Fig. 2.1) is obtained for many reactions. In such cases, the

slope of the line is $-E_a/2.303R$ and the intercept as $\frac{1}{T} = 0$ gives $\log A$.

Eqs. 2.59 is also written in the exponential form as follows:




Fig. 2.1. Plot of $\log k$ against $\frac{1}{T}$.

$$k = A e^{-E_a/RT} \quad \dots(2.61)$$

A possible reason for the deviation from Arrhenius equation in some reactions is that A and E_a may vary with temperature. The temperature dependence of Arrhenius factor will be discussed in collision theory. In the present discussion, we consider that A and E_a are constant for a reaction. If the activation energy is high for a reaction, it means that the temperature dependence of the reaction rate is also high. In such cases, even a small change of temperature results in a large change in the rate constant.

Although activation energy of a reaction can be calculated from $\log k$ vs $1/T$ plot, another way of obtaining it is to calculate rate constants (k_1 and k_2) at two temperatures (T_1 and T_2). Assuming E_a and A to be constant and using Eq. 2.60, we get.

$$\log k_1 = \log A - E_a/2.303RT_1 \quad \dots(2.62)$$

and $\log k_2 = \log A - E_a/2.303 RT_2 \quad \dots(2.63)$

Subtracting the terms in Eq. 2.63 from those in Eq. 2.62,

$$\begin{aligned} \log \frac{k_2}{k_1} &= \frac{-E_a}{2.303R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right) \\ &= \frac{-E_a}{2.303R} \frac{(T_2 - T_1)}{T_1 T_2} \\ \text{i.e., } \log \frac{k_2}{k_1} &= \frac{E_a}{2.303R} \frac{(T_2 - T_1)}{T_1 T_2} \quad \dots(2.64) \end{aligned}$$

The unit of A depends on the unit of k . For first order reactions, A has s^{-1} unit which is the same as the unit for frequency. This could be a reason for its name, frequency factor. A is also called the pre-exponential factor since it precedes the exponential term in Eq. 2.61

Let us work out an example.

Example 4

The rate constants for the decomposition of SO_2Cl_2 are $1.01 \times 10^{-6} \text{ s}^{-1}$ at 552 K and $3.85 \times 10^{-5} \text{ s}^{-1}$ at 600 K. Calculate the activation energy and the frequency factor for the assuming them to be independent of temperature.

Solution

From Eq. 2.64,

$$\begin{aligned} E_a &= 2.303 \frac{RT_1 T_2}{(T_2 - T_1)} \log \frac{k_2}{k_1} \\ E_a &= \left(\frac{2.303 \times 8.314 \times 552 \times 600}{48} \log \frac{3.85 \times 10^{-5}}{1.01 \times 10^{-6}} \right) \text{ J mol}^{-1} \end{aligned}$$

Substituting for $T = 600 \text{ K}$, $E_a = 2.09 \times 10^5 \text{ J mol}^{-1}$ and $k = 3.85 \times 10^{-5} \text{ s}^{-1}$ in Eq. 2.60

$$\begin{aligned} \text{we get, } \log A &= \log k + E_a/2.303RT \\ &= -4.4145 + \frac{2.09 \times 10^5}{2.303 \times 8.314 \times 600} \end{aligned}$$

$$\log A = -4.4145 + 18.1924 = 13.7779$$

$$A = \text{Antilog of } 13.7779$$

$$A = 6.00 \times 10^{13} \text{ s}^{-1}$$

We can understand the significance of the terms, E_a , A and $e^{-E_a/RT}$ during the discussion on collision theory of reaction rates.

Collision Theory

Collision theory is applicable to bimolecular reactions in gas phase. With some modifications, this can be applied to unimolecular (see Sec. 18.9) and termolecular reactions also. We explain collision theory using a gas-phase bimolecular elementary reaction of the following.



As per collision theory, the rate of a bimolecular reaction depends on

- The total collision frequency and
- Boltzmann factor.

Although the steric factor is also to be considered while calculating the reaction rate, it will be treated under the refinement of collision theory.

To understand the significance of Eq. 2.26, see Appendix I.

$$\text{average speed of a gas molecule } (\bar{u}) = \left(\frac{8RT}{\pi M_m} \right)^{1/2}$$

$$\text{i.e., Average speed of a gas molecule } (\bar{u}) = \left(\frac{8k_b T}{\pi m} \right)^{1/2}$$

$$\text{Since } R = N_A k_b, M_m = N_A m$$

Where M_m is the molar mass and m is the mass of one molecule of gas and N_A is Avogadro constant.

While calculating the relative motion of particles, it is customary to use reduced mass in the place of mass of one molecule of the gas.

$$\text{Hence average relative Speed of the molecules of X and Y} = \left(\frac{8k_b T}{\pi \mu} \right)^{1/2}$$

which is same as Eq. . . . 2.68

$$\mu = \frac{M_x M_y}{M_x + M_y} \frac{1}{N_A} \quad \dots 2.72$$

where M_x , M_y and N_A are the molar mass of X molar mass of Y and Avogadro constant

Eq. 2.72 comes from Eq. 2.70 since,

Mass of one molecule of a substance

$$= \frac{\text{Molar mass}}{\text{Avogadro constant}}$$

Total Collision Frequency

Total collision frequency (Z_{XY}) is the number of collision between the molecules of X and the molecule of Y in unit time in unit volume. Only X-Y collision are counted but not X-X or Y –Y collisions, since only X-Y collisions are possible for the reaction indicated in Eq. 2.65

The total collision frequency (Z) in general can be derived using the following relationship:

$$Z = \begin{cases} \pi \times (\text{collision diameter})^2 \\ \times (\text{average relative speed of gas molecules}) \\ \times (\text{number density}) \\ \times (\text{number density}) \\ \times (\text{correction factor}) \end{cases} \dots(2.66)$$

For calculating the total collision frequency (Z_{XY}) among the molecules of X and Y, as per Eq. 2.66, we use the following relationship:

$$1) \quad \text{Collision diameter} = \sigma_{XY} = \frac{1}{2} (\sigma_X + \sigma_Y) \dots(2.67)$$

where σ_X and σ_Y are the diameters of the molecules, X and Y, respectively. The collision diameter σ_{XY} is the distance of closest approach between a molecule of X and a molecule of Y.

$$\text{ii) Average relative speed of the molecules of X and Y} = \left(\frac{8 k_b T}{\pi \mu} \right)^{1/2} \dots(2.68)$$

where k_b is the Boltzmann constant (subscript b is added to k to differentiate it from the rate constant), T is temperature and μ is reduced mass.

$$\text{Note that } \frac{1}{\mu} = \frac{1}{m_X} + \frac{1}{m_Y} \dots (2.69)$$

$$\text{or } \mu = \frac{m_X m_Y}{m_X + m_Y} \dots(2.70)$$

where m_X and m_Y are the masses of one molecule of X and Y, representatively.

iii) Let us now calculate the factor, (number density) x (number density). Since we have two types of molecules, X and Y, we have to consider number densities of both X and Y. Using Eq. 2.75

To understand the significance of Eq 2.66, see Appendix I.

$$\begin{aligned} \text{Average speed of} & & & = & \left(\frac{8RT}{\pi M_m} \right)^{1/2} \\ \text{a gas molecule (u)} & & & & \\ \text{i.e., Average speed of} & & & = & \left(\frac{8k_b T}{\pi m} \right)^{1/2} \\ \text{a gas molecule (u)} & & & & \\ & & & & \dots (2.71) \end{aligned}$$

Since $R = N_A k_b$, $M_m = N_A m$

Where M_m is the molar mass and m is the mass of one molecule of gas and N_A is Avogadro constant.

While calculating the relative motion of particles, it is customary to use reduced mass in the place of mass of one molecule of the gas.

$$\left[\begin{array}{l} \text{Hence average relative} \\ \text{speed of the molecules} \\ \text{of X and Y} \end{array} \right] = \left(\frac{8k_b T}{\pi \mu} \right)^{1/2}$$

this is same as Eq. (2.68)

$$\mu = \frac{M_X M_Y}{M_X + M_Y} \cdot \frac{1}{N_A} \dots (2.72)$$

where M_X , M_Y and N_A are the molar mass of X, molar mass of Y, and Avogadro constant.

Eq. 2.72 comes from Eq. 2.70

Since,

mass of one molecules of a substance

$$= \frac{\text{Molar mass}}{\text{Avogadro constant}}$$

Number density of a gas has been defined in Subsec. 2.28 of Unit 2.

Number density

$$\begin{aligned} & \frac{\text{Number of molecules of the gas}}{\text{Volume of the gas}} \\ &= \frac{\text{Pressure of the gas}}{\text{Boltzmann constant} \times \text{temperature}} \\ &= \frac{P}{k_b T} \quad \dots (2.73) \end{aligned}$$

Also note that

Concentration

$$\begin{aligned} &= \frac{\text{Number of moles of the gas}}{\text{Volume of the gas}} \\ &= \frac{P}{RT} \end{aligned}$$

i.e., Concentration of a gas

$$= \frac{P}{N_A k_b T} = \frac{\text{Number density}}{N_A} \quad \dots (2.74)$$

Where N_A is Avogadro constant.

$$\text{i.e., Number density of a gas} = N_A \times (\text{concentration of the gas}) \quad \dots (2.75)$$

Boltzmann factor is helpful in calculating the number of molecules possessing energy equal to or greater than a particular value. Boltzmann factor, in other words, is helpful in calculating the population of the energy levels.

$$[X] = \frac{\text{Number of moles of X}}{\text{Volume}}$$

$$= \frac{\text{Number of molecules of X}}{N_A \times \text{Volume}}$$

$$= \frac{\text{Number density of X}}{N_A}$$

(number density of X) × (number density of Y)

In the case of collision between the molecules of X and Y (i.e., between molecules of different gases), there is no necessity for the correction factor. It is because we calculate the collisions between; each molecule of X and each molecule of Y. Each collision is counted only once. So omitting the correction factor and using Eqs. 2.66, 2.68 and 2.79, we get,

$$Z_{XY} = \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A^2 [X][Y] \quad \dots(2.79)$$

Thus we have obtained a relationship useful in calculating the total collision frequency for the collision between each molecule of X and each molecule of Y. Next we study the significance of Boltzmann factor.

Boltzmann Factor

You must realise that not all collisions between the molecules of X and Y would result in the product formation. Only those collisions, in which, the energy of the colliding molecules equals or exceeds some critical value E_a (known as activation energy as per Arrhenius equation), are effective in bringing about the reaction between X and Y. If $E_a \gg RT$, then the Boltzmann factor, $e^{-E_a/RT}$ gives the fraction of the collisions in which the colliding molecules possess energy equal to or greater than the activation energy.

$$\text{Boltzmann factor} = e^{-E_a/RT} \quad \dots(2.80)$$

Calculation of Reaction Rate

The product of the total collision frequency and the Boltzmann factor gives the **number of molecules of X or Y in unit volume reacting per unit time**. This follows from the definitions of the terms, total collision frequency and the Boltzmann factor. In order to obtain the reaction rate in terms of concentrations of X or Y (or the **number of moles of X or Y** unit volume) consumed per unit time, we have to divide the product, $Z_{XY} e^{-E_a/RT}$ by Avogadro constant;

$$\begin{aligned} \text{Reaction rate} &= \frac{-d[X]}{dt} = \frac{-d[Y]}{dt} \\ &= \frac{Z_{XY} e^{-E_a/RT}}{N_A} \quad \dots(2.81) \end{aligned}$$

Using Eqs. 2.79 and 2.81,

$$\text{reaction rate} = \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A^2 [X][Y] e^{-E_a/RT} \times \frac{1}{N_A}$$

$$= \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A [X] [Y] e^{-E_a/RT} \quad \dots(2.82)$$

By definition, the reaction rate for a bimolecular elementary reaction as per Eq.2.65 is as follows;

$$\text{Reaction rate} = k [X] [Y] \quad \dots (2.83)$$

Comparing Eqs. 2.82 and 2.83

$$k = \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A [X][Y] N_A e^{-E_a/RT} \quad \dots (2.84)$$

Eq. 18.155 gives the theoretical value of the rate constant for a bimolecular reaction as per collision theory, $\pi \sigma_{XY}^2$ is called the mean collision cross-section.

Collision Theory and Arrhenius Theory – a Comparison

You compare Arrhenius Equation (Eq. 2.61) with Eq. 2.84. You can see that the frequency factor A is given by,

$$A = \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A \quad \dots (2.85)$$

From Eq. 2.85, you can see that A is not independent of temperature as predicted by Arrhenius equation. However, over a short range of temperature, the variation in A is not significant. Arrhenius factor can be written as a product of A' and $T^{1/2}$ where A' is a temperature-independent constant and T is temperature.

$$A = A' T^{1/2}$$

Where
$$A' = \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A$$

The factor $T^{1/2}$ indicated the temperature dependence of A as given in Eq. 2.85

$$\text{Hence } \log A = \log A' T^{1/2} = \log A' + 1/2 \log T$$

Using this in Eq. 2.60

$$\log k = \log A' + 1/2 \log T - E_a/2.303 RT$$

$$\text{or } \log k - 1/2 \log T = \log A' - E_a/2.303 RT$$

A better method of arriving at the experimental values of E_a and A is to plot $\log k - 1/2 \log T$ against $1/T$.

The slope of the plot gives E_a value since

$$\text{Slope} = -E_a/2.303 R$$

The intercept gives $\log A'$ value from A' values, A at any temperature can be found out since

$$A = A' T^{1/2}$$

The value of A so obtained is the experimental value.

It was found that in the case of reaction between simple molecules, the agreement between the experimental value of A and the value obtained as per collision theory (Eq. 2.85) is fairly good. In the case of reactions involving complex molecules, there is a discrepancy between the two values of A . To explain the discrepancy between the two values of A , a refinement was suggested for Eqs. 2.84 and 2.85 in terms of steric factor.

Steric Factor

Although the molecular collision may have requisite energy, the reaction would take place only if the molecules have proper orientation. In other words, the reacting species must have proper spatial orientation for the reaction to occur. In the case of complex molecules, the probability of attaining proper orientation for the reaction is much less as compared to simple molecules. To stress the need for the poor spatial requirement, **steric factor or probability factor (P)** was also added to the right hand side of Eq. 2.84

$$\text{Hence } k = P \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A N_A e^{-E_a/RT} \quad \dots (2.86)$$

$$A = P \pi \sigma_{XY}^2 \left(\frac{8k_b T}{\pi \mu} \right)^{1/2} N_A \quad \dots (2.87)$$



$$\text{Also, } P = \frac{A_{\text{Experimental}}}{A_{\text{Theoretical}}} \quad \dots (2.88)$$

The steric factor is smaller for reaction between complex molecules. We expect the steric factor to be less than unity. But many fast reactions are known for which the steric factor is much greater than unity Collision theory. Collision theory

cannot explain such cases. Let us now study the **Activated Complex Theory** which gives a better method of calculating reaction rates.


Activated Complex Theory

The activated complex theory or the absolute theory of reaction rates depicts the formation of activated complex (A^\ddagger) from the reactants (X and Y) as a preceding step for the formation of the product, P.



The main features of the activated complex theory are given here.

The reactant molecules come into contact with each other. In this process, a few bonds get distorted, some bonds start forming with the exchange or release of atomic or groups. The composite molecule so formed from the reactants prior to the formation of the product is called the activated complex. The activated then decomposes to give the product. The reaction sequence could be represented as in Fig 2.2



Activated complex is the configuration of the atoms which the reactant molecules have near the top of the energy barrier that separates the reactants from the products. Transition state is the highest point in the potential energy curve.

Fig. 2.2. Graphical representation of the change in potential energy as a function of reaction coordinates.


The total potential energy of the system is depended in the y-axis and the **reaction coordinate** in the x-axis. Reaction coordinate is the sequence of simultaneous changes in bond distances and bond angles. Such changes result during the formation of the products from the reactants.

Consider the reaction between a molecule of H_2 and a molecule of I_2 . To start with, let us imagine that the two molecules are far part and the total potential energy of the system is the sum of the potential energies of H_2 and I_2 . This part of the reaction course is represented by the horizontal portion AB of the curve in

Fig. 18.11. As the two molecules approach each other to such an extent that the orbitals begin to overlap (point B in the curve), H–H and I–I bonds begin to stretch and H–I bond begins to form. The total potential energy starts increasing and this is represented by raising portion of the curve BC. As the extent of H–H and I–I bond breakage and H–I bond formation increase, a point is reached when the potential energy is maximum (point C). The activated complex, represented below as a composite molecule, has the maximum potential energy.

Activated complex is the configuration of the atoms which the reactant molecules have near the top of the energy barrier that separates the reactants from the products. Transition state is the highest point in the potential energy curve.

The bond-breaking is an energy demanding process and the bond-making is an energy releasing process. The net energy requirement for the formation of the activated complex and its decomposition to products must be available through the translational or the vibrational energy of the reactants.



Activated Complex

The maxima point in the potential energy curve is called the transition state. Even a slight distortion of the bonds in the form of compression of H – I bond and stretching of H – H and I – I bonds enables the activated complex pass through the transition state. The path along CD represents the course, of the events which result in the complete breakage of H – H and I – I bonds along with the formation of H – I bond. The horizontal portion DE represents the total potential energy of two H – I molecules. Although a fraction of the activated complex molecules could form the reactants (along the path CB) the formation of the products is almost a certainty, once the activated complex is at the transition state. The fraction of the activated complex converted into products is called the transmission coefficient and, in majority of cases, it is unity.


Energy Requirement for the Reaction

Now let us consider the energy criteria for the reaction. The energy requirement for the reactants to cross the energy barrier is to be met from translational or the vibrational energy of the molecules. At the transition state, the activated complex has some complicated vibration like motion of all atoms. The activated complex has one particular mode of vibration along which it is unstable. If the activated

complex vibrates with the frequency corresponding to this vibrational mode, the activated complex decomposes into products.

Rate Constant Calculation using Activated Complex Theory

Based on statistical thermodynamics, Eyring developed the activated complex theory. The basic postulate of the theory is that there exists an equilibrium between the activated complex and the reactants. Let us consider the **bimolecular gas phase reaction**,



where X and Y are the reactants, and A^\ddagger is the activated complex: The activated complex then decomposes to give the product, P



The rate of formation of the product depends on,

- i) the concentration of the activated complex and
- ii) the frequency which it is converted into the product. This is the frequency of one of the vibrational modes with respect to which the activated complex is unstable. Using detailed calculations it is possible to derive an expression useful in calculating the rate constant (k) the elementary reaction, $X + Y \rightarrow P$ (for which the steps are given in Eqs. 2.89 and 2.90). We shall only state the final expression without going through the derivation in full.

$$k = \frac{RT}{P^0} \cdot \frac{k_b T}{p^0} k_p = \frac{Rk_b T^2}{p^0 h} \cdot k_p \quad \dots (2.91)$$

where $K_p =$ The Equilibrium constant for the formation of the activated complex after adjusting for its vibration with respect to which it is unstable.

$T =$ Temperature

$k_b =$ Boltzmann constant

RT / p^0 is the correction term where p^0 is the standard pressure (1bar)

$h =$ Planck constant

$R =$ Gas constant

Using van't Hoff isotherm ... (2.92)

$$\Delta G^\ddagger = -RT \ln K_p$$

where ΔG^\ddagger is molar Gibbs energy of activation. ... (2.93)

Hence, $K_p = e^{-\Delta G^\ddagger/RT}$

Using Eq. 2.93 in Eq. 2.91

$$k = \frac{Rk_b T^2}{hp^0} e^{-\Delta G^\ddagger/RT} \quad \dots (2.94)$$

Using

$$\Delta G^\ddagger = \Delta H^\ddagger - T\Delta S^\ddagger \quad \dots (2.95)$$

where ΔH^\ddagger and ΔS^\ddagger are the molar enthalpy of activation and entropy of activation, respectively.

Using Eq. 2.95 and 2.94

$$k = \frac{Rk_b T^2}{hp^0} e^{-(\Delta H^\ddagger - T\Delta S^\ddagger)/RT}$$

$$k = \frac{Rk_b T^2}{hp^0} e^{-(\Delta H^\ddagger - RT\Delta S^\ddagger)/RT} \quad \dots (2.96)$$

Using differential form of Eq. 2.59, it is possible to find the following relationship for a bimolecular gas phase reaction (see Appendix II):

$$\Delta H^\ddagger = E_a - 2RT \quad \dots(2.97)$$

Using Eq.s 2.96 and 2.97

$$k = \frac{Rk_b T^2}{hp^0} e^{-(E_a - 2RT)/RT} e^{\Delta S^\ddagger/R}$$

$$k = \frac{Rk_b T^2}{hp^0} e^{-E_a/RT} e^2 e^{\Delta S^\ddagger/R} \quad \dots(2.98)$$

Comparing Arrhenius equation Eq 2.61 and with Eq. 2.98

$$A = \frac{Rk_b T^2}{hp^0} e^2 e^{\Delta S^\ddagger/R} \quad \dots (2.99)$$

$$\text{Hence } e^{\Delta S^\ddagger/R} = \frac{hp^0}{Rk_bT^2} e^2 \cdot A$$

$$\text{i.e., } e^{\Delta S^\ddagger/R} = 7.8119 \times 10^{-11} \frac{A}{T^2} \quad \dots (2.100)$$

(substituting for the constants)

Taking natural logarithms,

$$\Delta S^\ddagger/R = \ln 7.8119 \times 10^{-11} \frac{A}{T^2}$$

$$\begin{aligned} \Delta S^\ddagger &= 2.303R (\log 7.8119 \times 10^{-11} + \log A/T^2) \\ &= (19.15 \log A/T^2 - 193.6) \text{ J mol}^{-1} \text{ K}^{-1} \quad \dots (2.101) \end{aligned}$$

Thus ΔH^\ddagger , ΔS^\ddagger and ΔG^\ddagger can be calculated at any given temperature. If A and E_a are known from Arrhenius plot (Fig. 2.1).

While discussing collision theory or the activated complex theory, we consider only the gas-phase reactions. The study of reaction rates in solutions is complicated due to the role of the solvent. In this course, we shall not discuss the reaction rates in solutions.

To account for the discrepancy between the experimentally obtained value of A and the value calculated as per collision theory, we had to introduce steric factor in Eq. 18.157. The activated complex theory has the factor, $e^{\Delta S^\ddagger/R}$ which takes care of the steric factor automatically.

It is observed that ΔS^\ddagger is negative for many reactions. Such a negative Value indicates decrease in disorderliness. This is understandable since during collisions, the particles have to approach each other, thereby causing a decrease in randomness. The negative value of ΔS^\ddagger brings down the value of $e^{\Delta S^\ddagger/R}$ and hence of A as per Eq. 18.-170. A large negative value for the entropy of activation is generally observed for reactions involving complex molecules, the demand for proper orientation is more in the case of a complex molecule and this causes larger reduction in randomness. Thus the entropy of activation, and hence the frequency factor could be expected to be much less for reactions involving complex molecules.

SAQ 3

The second order rate constants of a reaction are given below at two temperatures:


T/K	298	308
$10^5 \times k / \text{M}^{-1} \text{s}^{-1}$	8.8	28

Calculate the activation energy of the reaction.

.....
.....
.....

SAQ 4

For the reaction,



Arrhenius factor at 298 k is $9.4 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$, Calculate entropy of activation for this reaction at 298 k.


.....
.....
.....

3.5 Complex Reactions

In the preceding units we have discoursed on the rate of chemical reactions and factors influencing it. Emphases were made on the rate laws of first and second order reactions which are the simple reactions. Not all chemical reactions proceed to a stage at which the concentrations of the reactions becomes vanishingly small. Such a reaction proceeds via a more complex reaction mechanism. Most industrial chemical reactions, the type likely to be encountered in a chemical laboratory or plant, involve multiple steps between the reaction and products. In this section we will consider increasingly complex reaction schemes, and see how to construct and solve the corresponding rate laws.

3.51 Parallel Reaction

It is not uncommon for reaction to produce more than one product, and the reaction is often kinetic and thermodynamic. The sequence is



$$\text{Rate} = \frac{d[A]}{dt} = -K_1[A] - K_2[A]$$

$$[A] = ae^{-(K_1+K_2)t}$$

The rate of the formation of the products are

$$\frac{d[B]}{dt} = K_1[A]$$

$$= K_1ae^{-(K_1+K_2)t}$$

and

$$\frac{d[C]}{dt} = K_2[A]$$

$$= K_2ae^{-(K_1+K_2)t}$$

Integrating the equation gives

$$[B] = \frac{K_1a}{K_1 + K_2} (1 - e^{-(K_1+K_2)t})$$

and


$$[C] = \frac{K_2a}{K_1 + K_2} (1 - e^{-(K_1+K_2)t})$$

The rate of the products formed is proportional to their rate constant

$$\frac{[B]}{[C]} = \frac{K_1}{K_2}$$

3.52 Reactions Approaching Equilibrium

Consider a reaction in which both forward and reverse reaction are first order as given by the scheme below.



The rate of change of [A] has two contributions. It is depleted by the forward reaction at a rate $K[A]$ but is replenished by the reverse reaction at a rate $K^1[B]$.

The net rate of change is therefore

$$\frac{d[A]}{dt} = -K[A] + K^1[B]$$

If the critical concentration of A and [B]_o is [A]_o and there is no B present initially, at all times $[A] + [B] = [A]_o$, and so

$$\begin{aligned} \frac{d[A]}{dt} &= -K[A] + K^1([A]_o - [A]) \\ &= -(K + K^1)[A] + K^1[A]_o \end{aligned}$$

The solution of this first order differential equation, with the critical condition [A]

= [A]_o is

$$[A]_t = [A]_o \left[\frac{K^1 + K e^{-(K+K^1)t}}{K_1 + K^1} \right]$$

When time t approaches infinity the concentrations reach their equilibrium values

$$[A]_\alpha = K^1[A]_o / (K_1 + K^1) \text{ and}$$


$$[B]_\alpha = [A]_o - [A] = K[A]_o / (K + K_1)$$

The ratio of these equilibrium concentrations, which is the equilibrium constant is

$$K_c = \frac{[B]_{\alpha}}{[A]_{\alpha}} = \frac{K}{K^1}$$

3.53 Other Types of Equilibria

In the case of a reaction that is bimolecular and second-order in both directions as shown by the scheme below



The rate of change of the concentration of A, as a result of the forward, and reverse reactions are



and



At equilibrium the net rate of change is zero. Hence, at equilibrium

$$-K[A] [B] + K^1 [C] [D] = 0, \text{ and so}$$

$$K_c = \left\{ \frac{[C][D]}{[A][B]} \right\}_{eq} = \frac{K}{K^1}$$

In the case of a reaction which proceeds by a sequence of simple reactions, such as



$$v_A \text{ reverse} = K^1_a [C] [D]$$




$$v_C \text{ reverse} = K^1_b [E] [F]$$

at equilibrium all the reactions are individually at equilibrium, so that,

$$\left\{ \frac{[C][D]}{[A][B]} \right\}_{eq} = \frac{K_a}{K_a^1} \text{ and } \left\{ \frac{[E][F]}{[C]} \right\}_{eq} = \frac{K_b}{K_b^1}$$

The overall reaction equilibrium is



$$K = \left\{ \frac{[D][E][F]}{[A][B]} \right\}_{eq} = \left\{ \frac{[C][D][E][F]}{[A][B][C]} \right\}_{eq}$$

$$= \left\{ \frac{[C][D]}{[A][B]} \right\}_{eq} \left\{ \frac{[E][F]}{[C]} \right\}_{eq} = \frac{K_a K_b}{K_a^1 K_b^1}$$


When the overall reaction is the sum of a sequence of steps

$$K = \frac{K_a K_b}{K_a^1 K_b^1}$$


Where K_∂ are the rate constants for the individual steps and K^1 refers to the corresponding reverse reaction steps.

3.54 Consecutive Reactions

Some reactions proceed through the formation of an intermediate as in radioactive decay



Consider a first-order consecutive reaction as given below



The rate of disappearance of A is

$$\frac{d[A]}{dt} = -K[A]$$

and that of formation of B and C are

$$\frac{d[B]}{dt} = K_1[A] - K_2[B]$$

$$\frac{d[C]}{dt} = K_2[B]$$

At initial time $t = 0$, concentration of $[A] = [A]_0$ and those of $[B] = 0$ and $[C] = 0$

The first equation is a first order rate law, and so $[A] = [A]_0 e^{-K_1 t}$

The rate of second equation is

$$\frac{d[B]}{dt} = K_1[A]_0 e^{-K_1 t} - K_2[B]$$


$$\frac{d[B]}{dt} = K_2[B] = K_1[A]_0 e^{-K_1 t}$$

By integrating the equation, the solution is

$$[B] = \frac{K_1}{K_2 - K_1} [A]_0 \left\{ 1 + \left(\frac{e^{-K_1 t} - K_2 e^{-K_2 t}}{K_2 - K_1} \right) \right\}$$

3.55 Pre-equilibria

In this case, a consecutive reaction in which an intermediate reaches an equilibrium with the reactions before forming a product, as given in the scheme below



Since we assume that A, B and (AB) are in equilibrium, we can write

$$K = \left\{ \frac{[(AB)]}{[A][B]} \right\}_{eq}$$

$$\text{with } K = \frac{K_1}{K_2}$$


By ignoring the fact that [AB] is slowly leaking away as it form C. The rate of formation of C may now be written as

$$\begin{aligned} \frac{d[C]}{dt} &= K_2[(AB)] \\ &= K_2 K_1 [A][B] \\ &= K [A][B] \end{aligned}$$

$$\text{Where } K = \frac{K_1 K_2}{K_2}$$

3.56 Enzyme Reaction

Another example of a pre-equilibrium reaction is the Michaelis-Menten mechanism of enzyme action. The proposed mechanism is



$$\frac{d(p)}{dt} + K_3[(ES)]$$

(ES) denotes a bound state of the enzyme E and its substrates S. In order to relate [(es)] to the enzyme concentration we write its rate law and then impose the steady-state approximation

$$\frac{d[(ES)]}{dt} = K_1[E][S] - K_2[(ES)] - K_3[(ES)] = 0$$

This solve to

$$[(ES)] = \left\{ \frac{K_1}{(K_2 + K_3)} \right\} [E][S]$$

[E] and [S] are concentration of the enzyme and substrate and $[E]_o$ in total concentration of enzyme

$$[E] + [(ES)] = [E]_o, \text{ a constant}$$

since only few E is added, we can ignore the fact that [S] differs slightly from [S] total.

Therefore

$$[(ES)] = \left\{ \frac{K_1}{K_2 + K_3} \right\} \{ [E]_o - [(ES)] \} [S]$$

Which rearranges to

$$[ES] = \frac{K_1 [E]_o [S]}{K_3 + K_2 + K_1 [S]}$$

It follows that the rate of formation of products is

$$\begin{aligned} \frac{d[P]}{dt} &= \frac{K_3 K_1 [E]_o [S]}{K_3 + K_2 + K_1 [S]} \\ &= \frac{K_3 [E]_o [S]}{K_m + [S]} \end{aligned}$$

where k_m is the Michael constant is

$$K_m = \frac{(K_3 + K_2)}{K_1}$$

3.57 Unimolecular Reaction

A number of gas phase reactions follows first-order kinesis and are believed to proceed through a unimolecular rate – determining stage. These are called unimolecular reactions. In the Lindemann-Heinshelwood mechanism it is supposed that a reactant molecule A collides with another M, a diluents gas

molecule, and becomes energetically excited at the expense of M's translational kinetic energy,



and the energised molecule might lose its excess energy by colliding with another



or the excited molecule might shake itself apart and form product



$$\frac{d[A^*]}{dt} = -K_3[A^*]$$

By applying steady state approximation to the net rate of formation of A^* ,

$$\frac{d[A^*]}{dt} = K_1[A][M] - K_2[A^*][M] - K_3[A^*] = 0$$

This solves to

$$[A^*] = \frac{K_1[A][M]}{K_3 + K_2[M]}$$

and so the rate law for the formation of P is

$$\frac{d[P]}{dt} = K_3[A^*] = \frac{K_1K_3[A][M]}{K_3 + K_2[M]}$$

If the rate of deactivation by A^* , M collisions is much greater than the rate of unimolecular decay, so that

$$K_2[A^*][M] \gg K_3[A^*] \text{ or } K_2[M] \gg K_3$$

Then we neglect K_3 in the denominator and obtain

$$\frac{d[P]}{dt} = \frac{K_1K_3[A][M]}{K_2[M]} = \left(\frac{K_1K_3}{K_2} \right) [A]$$

a first-order rate law, as we set out to show.

The Lindemann-Hinshelwood Mechanism can be tested because it predicts that as the concentration of M is reduced, the reaction should switch to overall second-order kinetics. This is because when $K_2[M] \ll K_3$, the rate is approximately

$$\frac{d[P]}{dt} = \frac{K_1 K_3 [M][A]}{K_3} = K_1 [A][M]$$

The physical reason for the change of order is that at low pressure the rate-determining step is the bimolecular formation of $[A^*]$, if we write the full rate law as

$$\frac{d[P]}{dt} = K_{eff}[A]$$

$$K_{eff} = \frac{K_1 K_3 [M]}{K_3 + K_2 [M]}$$

then the expression for the effective rate constant can be rearranged to

$$\frac{1}{K_{eff}} = \frac{1}{K_1 [M] + \frac{K_2}{K_1 K_3}}$$

SAQ 5


The rate law for the reaction $2A+B \rightarrow X + 3D$ was expressed as $\text{rate} = K [A][B]$, what are the units of K? express the rate law in terms of the rate of change of concentration of (a) A, (b) c.

3.6 Summary

In this unit we discussed the steps followed in arriving at the mechanisms of simple reactions. We described the theory of unimolecular reaction rates. Arrhenius equation, collision theory and the activated complex were discussed.

3.7 Tutor mark assessment questions.

- 1) In the reaction




the mechanism proposed below agrees with the experimentally observed rate law:



What is the rate law?

- 2) For the decomposition of N_2O at 773 K, $k = 1.00 \times 10^{-5} \text{ s}^{-1}$ and $E_a = 250 \text{ KJ mol}^{-1}$. Calculate A using Arrhenius equation.
- 3) For the reaction,



the steric factor is 1.7×10^{-6} and mean collision cross-section ($\pi \sigma^2$) is 0.46 nm^2 . Calculate A at 628 K. Use Eq. 2.87

Given: i) $k_b = 1.381 \times 10^{-23} \text{ J. K}^{-1}$

ii) The relative molecular masses of ethylene and hydrogen are 28.05 and 2.016, respectively.

iii) $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

- 4) Explain the reason for the negative value of entropy of activation in the case of many reactions.

3.8 Answers

Self Assessment Questions.

- 1
- i) Rate = $k[\text{O}_3][\text{NO}]$
 - ii) Rate = $k [\text{CH}_3\text{NC}]$
 - iii) Rate = $k[\text{Cl}_2]$

2. The rate law is given below as per the proposed mechanism:

$$\text{Rate} = k_3[\text{NO}_3] [\text{NO}]$$

$$\text{But } K = \frac{k_1}{k_2} = \frac{[\text{NO}_3]}{[\text{NO}][\text{O}_2]}$$

$$\text{Hence, } [\text{NO}_3] = K [\text{NO}] [\text{O}_2]$$

Using (2) in (1),

$$\begin{aligned} \text{rate} &= k_3 K [\text{NO}]^2 [\text{O}_2] \\ &= k [\text{NO}]^2 [\text{O}_2] \end{aligned}$$

this is same as the rate law obtained experimentally

3) Using Eq.2.64

$$\begin{aligned} E_a &= \frac{2.303 RT_1 T_2}{(T_2 - T_1)} \log k_2/k_1 \\ &= \frac{2.303 \times 8.314 \times 298 \times 308}{10} \log \frac{28 \times 10^{-5}}{8.8 \times 10^{-5}} \\ &= 88.3 \text{ kJ mol}^{-1} \end{aligned}$$

4) According to Eq. 2.101

$$\begin{aligned} \Delta S^\ddagger &= (19.15 \log A/T^2 - 193.6) \text{ J mol}^{-1} \text{ K}^{-1} \\ &= \left(19.15 \log \frac{9.4 \times 10^9}{298^2} - 193.6 \right) \text{ J mol}^{-1} \text{ K}^{-1} \\ \Delta S^\ddagger &= -97.4 \text{ J mol}^{-1} \text{ K}^{-1} \end{aligned}$$

5) Ans.D – K is express as $\text{M}^{-1}\text{S}^{-1}$

$$v_A = -K^1 [A] [B], K^1 = K/V$$

$$v_C = -K^{11} [A] [B], K^{11} = 3K/V$$

Tutor mark assessment Question

1) Rate = $k_1[\text{NO}_2\text{Cl}]$

2) Using Eq. 2.60 $\log A = E_a/2.303 RT + \log k$

$$= (250 \times 10^3 / 2.303 \times 8.314 \times 773) + \log (1.00 \times 10^{-5})$$

i.e. $\log A = -5.00 + 16.89$
 $A = 7.8 \times 10^{11} \text{ s}^{-1}$

3) Mass of one molecule of a substance = $\frac{\text{Molar mass}}{N_A}$

Mass of one molecule of ethylene = $\frac{28.05 \times 10^{-3}}{6.022 \times 10^{23}} \text{ kg}$

Mass of one molecule of hydrogen = $\frac{2.016 \times 10^{-3}}{6.022 \times 10^{23}}$

Using Eq. 2.70 $\mu = \frac{28.05 \times 2.016}{30.07 \times 6.022 \times 10^{26}} \text{ kg}$

$$= 3.12 \times 10^{-27} \text{ kg}$$

$$\left(\frac{8k_b T}{\pi \mu} \right)^{1/2} = \left(\frac{8 \times 1.381 \times 10^{-23} \times 628}{3.143 \times 3.12 \times 10^{-27}} \right)^{1/2} \left(\frac{\text{J K}^{-1} \text{ K}}{\text{kg}} \right)^{1/2}$$

$$= 2.66 \times 10^3 \text{ m s}^{-1}$$

Using Eq. 2.87

$$A = P \pi \sigma^2 \left(\frac{8 k_b T}{\pi \mu} \right)^{1/2} N_A$$

$$= 1.7 \times 10^{-6} \times 0.46 \times (10^{-9} \text{ m})^2 \times 2.66 \times 10^3 \text{ m s}^{-1} \times 6.022 \times 10^{23} \text{ mol}^{-1}$$

$$= 1.25 \times 10^3 \text{ m}^3 \text{ mol}^{-1} \text{ s}^{-1}$$

$$= 1.25 \times 10^6 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1} \quad \left. \vphantom{1.25 \times 10^6} \right\} (1 \text{ m}^3 = 10^3 \text{ dm}^3)$$

$$= 1.25 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$$

See under activated complex theory in sec. 2.4

UNIT 4 HETEROGENEOUS REACTIONS AND CATALYSIS

- 4.0 Introduction
 - Objectives
- 4.1 Surface Tension of Solutions
- 4.2 Adsorption on Solids
 - 4.21 Adsorption of Gases by Solids
- 4.3 Adsorption Isotherms
 - 4.3.1 Langmuir Adsorption Isotherm
- 4.4 Physisorption and Chemisorption
 - 4.41 Physisorption
 - 4.42 Chemisorption
- 4.5 Modern Methods of Surface Studies
- 4.6 Types of Catalysts
- 4.7 Mechanisms of Catalytic Reactions
- 4.8 Inhibition and Poisoning
- 4.9 Conclusion
- 4.10 Summary
- 4.11 Tutor Marked Assignment
- 4.12 References/Further Readings

4.0 INTRODUCTION

In Unit 18, we discussed the dependence of the rates of reactions on the concentration of the reactants and temperature. We explained the characteristics of the colloids which include surface activity also. In this unit, we shall discuss the effect of surface on chemical reactivity.

Surface, in the chemical sense, is a phase boundary. Geometrically a surface has an area but no thickness. Surface is an **interface region** where one phase ends and the other begins. Chemically it is a region in which the properties vary from one phase to another. The transition occurs over distance of molecular dimensions. Thus for a chemist, surface has a thickness which shrinks to zero in an ideal condition of a geometrical description. In this unit we shall discuss the properties of an interface which may be liquid/vapour, solid/liquid or solid/gas. We shall discuss adsorption of gases on solid surface. Such a discussion is important since many of the chemical reactions in industry or in the biological systems take place on the interface. We shall also explain the different types of catalysts, inhibition of catalysts and catalytic poisoning.

OBJECTIVES

After studying this unit, you should be able to:

- define an interface,
- state the relationship between the concentration of a surface active solute and the surface tension of a solution,
- discuss different isotherms for adsorption on solid surfaces,
- state the role of photoelectron spectroscopy in surface chemical analysis
- describe the functioning of different types of catalysts, and
- explain the inhibition of chemical reactions and poisoning of catalysts.

4.1 Surface Tension of Solutions

Liquids assume a shape with minimum surface area. The molecules in the bulk have less energy than the molecules in the surface, since a molecule in the bulk interacts with larger number of molecules as compared to a molecule on the surface. You are aware that any system tries to assume a state in which it has minimum energy. Hence, a liquid tries to take a shape which has the least surface area. The shape assumed is spherical, as a sphere has the smallest surface to volume ratio. The force that opposes the increase in area of a liquid is referred to as surface tension. In have defined surface tension as a force per unit length acting on the surface opposing the expansion of surface area. An alternative definition was also suggested that surface tension is the surface energy per unit area of the surface. Normally the surface tension values reported are for the liquid-vapour interface in presence of air.

When we dissolve a solute in a solvent, the surface tension of the solution changes. A relationship could be derived to establish the fact that the concentration of the solute that lowers the surface tension would tend to be more on the surface of a solvent as compared to that in the bulk. This is the basis of **Gibbs adsorption isotherm**, about which we do not intend giving a detailed discussion. Substances which produce a remarkable lowering of interfacial tension are called 'surface active agents or surfactants'. We have seen one of the applications of the surfactants in where we deal with the cleaning action of soaps and detergents. Another application of surfactants is in the formation of surface films. Some insoluble substances such as long-chain fatty acids and alcohols could spread on water surface to form a thin film. The formation of such surface films using long-chain alcohols has been helpful in retarding the evaporation of water from reservoirs.

SELF ASSESSMENT EXERCISE 1

What is the essential characteristic of the surface active agent?

.....
.....
.....

4.2 Adsorption on Solids

In Sec. 3.1, we studied that the molecules on the surface of a liquid experiences an inward pull. The liquid surface is thus in a state of unsaturation. The surface of a solid also behaves in a similar manner. In a solid, the ions or the molecules at the surface of a crystal do not have all their valencies satisfied by union with other particles. Such forces also arise due to the fact that when a new solid surface is created by breaking a solid, some interatomic bonds are broken and some of the valencies of surface atoms are left unsatisfied. As a result of these residual forces, the surface of the solid has a tendency to attract and retain available molecules and other particles towards itself; such a condition is helpful in decreasing the surface energy of a solid. The molecules so attracted are retained on the surface than in the bulk of the solid. This phenomenon of higher concentration of a substance on the surface of a solid is called **adsorption**. The substance attracted to the surface is called **adsorbate** while the substance to which it is attached is called **adsorbent**. For example, charcoal adsorbs acetic acid when kept in contact with it; here, acetic acid is adsorbate and charcoal is adsorbent.

Adsorption should be clearly distinguished from absorption. In absorption, the substance is not only retained on the surface but passes through the surface and is distributed throughout the bulk of the solid. Thus, anhydrous calcium chloride absorbs water to form a hydrate while acetic acid is adsorbed from its solution by charcoal. Sometimes the word 'sorption' is used when there is a doubt whether a process is true adsorption or absorption.

Note

Porous substance; a substance having tiny opening through which fluids or air could pass.

The extent of adsorption by charcoal can be increased by subjecting charcoal to a process of activation. It involves heating of wood charcoal between 625 K and 1275 K in vacuum, air, steam, chlorine or carbon dioxide. During activation, hydrocarbons and other impurities are removed from charcoal leading thereby to a large surface area for adsorption. The resulting substance is called activated charcoal.

4.21 Adsorption of Gases by Solids

The studies of the adsorption of gases by solids are similar to those of the adsorption of liquids by solids. In this unit, we shall mainly study the adsorption of gases by solids.

Although all solids adsorb gases, the effects are not evident unless the adsorbent is porous and has a very large area for a given mass. That is why silica gel and charcoal, which have porous structure, are very effective as adsorbing agents.

The amount of a gas adsorbed by a solid depends upon

- the nature of the gas and adsorbent,
- the surface area of the adsorbent, and
- temperature and pressure of the adsorbent-adsorbate system

Table 2.1 gives the volume of various gases adsorbed by 1.00 kg of charcoal at 288K. The volumes of gases have all been reduced to 273K and 1.013×10^5 Pa pressure. It can be seen from Table 2.1 that gases which can be liquefied easily are more readily adsorbed.

Table 2.1: Adsorption of Gases on Charcoal

Gas	Volume adsorbed m^3	Critical temperature k
H ₂	4.7×10^{-3}	33
N ₂	8.0×10^{-3}	126
CO	9.3×10^{-3}	134
CO ₂	4.8×10^{-2}	304
HCl	7.2×10^{-2}	324
H ₂ S	9.9×10^{-2}	373
NH ₃	1.8×10^{-1}	406

The total amount of the gas adsorbed increases with the surface area of the adsorbent. During adsorption an equilibrium is established between the gas in contact with the solid and the gas on the surface. An increase in temperature decreases the amount of the gas adsorbed.

In the text section, we shall discuss quantitative relationships regarding adsorption.

4.3 Adsorption Isotherms

The amount of a substance adsorbed by an adsorbent temperature depends upon the concentration or pressure of an adsorbate. For the adsorption of a substance (adsorbate) present in a solution by a solid adsorbent, Freundlich gave an empirical equation. This equation gives the relationship between the mass of the adsorbate (x) adsorbed by a particular mass (m) of the adsorbent and the equilibrium concentration of the adsorbate (c) in the solution at a particular temperature as given below:

$$\frac{x}{m} = Kc^{1/n} \quad \dots 2.1$$

where K and n are constants. Eq. 1.1 is a form of Freundlich adsorption isotherm. By plotting $\log \frac{x}{m}$ against $\log c$ (Fig. 1.1), we can evaluate K and n . The values of K and n depend on

- nature of the adsorbate,
- nature and particle size of the adsorbent, and
- temperature.

It is worth mentioning that as the particles size becomes smaller, the surface area increases enormously. This increases the adsorbing capacity of an adsorbent. Eq. 2.1 could be slightly modified to express the adsorption of a gas by a solid as given below:

$$\frac{x}{m} = Kp^{1/n} \quad \dots\dots 2.2$$

In this equation, p stands for the pressure of the gas adsorbate; other terms have the same significance as given in Eq. 2.1




Fig 2.1: Graphical representation of Freundlich adsorption isotherm.

Freundlich isotherm (Eq 2.1 or 2.2) is applicable only if the concentration or pressure of the adsorbate is low.

4.31 Langmuir Adsorption Isotherm

Langmuir obtained a relationship for the adsorption of a gas by a solid. Langmuir started with the assumptions stated below:

- The adsorbed gas behaves ideally in the vapour phase; there are no attractive or repulsive forces among the gas molecules.
- The surface of a solid is homogeneous and there are a fixed number of adsorption sites. Each site has the same attraction for the gas molecules.
- Each adsorption site can adsorb only one molecule. A solid surface cannot form a layer more than a single molecule in depth. In other words, the adsorption of a gas could lead only to the formation of unimolecular layer on the solid.
- There is an equilibrium between the condensation of gas molecules on the adsorbent and their desorption from it. The initial rate of condensation of

gas molecules on the solid surface is high and it decreases as the surface available for adsorption decreases. The desorption can occur by thermal agitation and the rate of desorption will depend upon the amount of solid surface covered by gas molecules. It will increase as the surface becomes more and more saturated. At a certain stage, the rates of condensation and desorption become equal and an equilibrium is established.

- Using the above assumptions, Langmuir equation for adsorption can be derived as follows:

Note:

Desorption is the process of release of the adsorbed molecules. Desorption may also be called the evaporation of the adsorbed molecules.

Let the fraction of the total surface covered by gas molecules be θ ; then the surface available for adsorption is $1 - \theta$. According to kinetic theory of gases, the rate at which the molecules strike the unit area of surface is proportional to the pressure of the gas. If p is the equilibrium gas pressure, then,

$$\text{rate of condensation} = a(1-\theta)p$$

$$\text{or rate of condensation} = k_1(1-\theta)p \quad \dots\dots 2.3$$

where k_1 is a constant of proportionality.

Rate of evaporation from the surface will be proportional only to the fraction of the surface which has adsorbed gas molecules on it. Thus,

$$\text{rate of evaporation} = k_2 \theta \quad \dots\dots 2.4$$

k_2 is also a proportionality constant.

At equilibrium, the rates of condensation and evaporation are equal. Hence,

$$K_1(1-\theta)p = k_2 \theta \quad \dots\dots 2.5$$

Rearranging Eq. 2.5, we can get,

$$\theta = \frac{k_1 p}{k_2 + k_1 p} = \frac{(k_1/k_2)p}{1 + (k_1/k_2)p} \quad (\text{Dividing the numerator and Denominator by } K_2)$$

$$\theta = \frac{Kp}{1 + Kp} \quad (\text{where } k_1/k_2 = K, \text{ another constant}) \quad \dots\dots\dots 2.6$$

Eq. 2.6 could be modified to find a relationship between the amount of gas adsorbed and the gas pressure. The amount of gas adsorbed (y) at a pressure p is proportional to the fraction of the total surface covered (θ) by the gas molecules.

i.e., $y \propto \theta$

Hence, $y = y_m \theta$ or $y/y_m = \theta$ 2.7

Where y_m is the proportionality constant and is equal to the amount of the gas molecules required to form a unimolecular layer; i.e., $y = y_m$ when $\theta = 1$

Using Eqs. 2.6 and 2.7

$$\frac{y}{y_m} = \frac{kp}{1 + kp}$$

$$y = \frac{y_m kp}{1 + kp}$$

or $\frac{p}{y} = \frac{1 + Kp}{y_m K} = \frac{1}{y_m K} + \frac{p}{y_m}$ 2.8

Eq. 2.8 is known as Langmuir adsorption isotherm.

When the gas pressure is low, p/y_m is small when compared to $\frac{1}{y_m k}$ hence $\frac{p}{y} =$

$$\frac{1}{y_m k} \text{ or } \frac{p}{y} = \text{constant, since } y_m \text{ and } K \text{ are constants.}$$

Or $p \propto y$

This means that at Low pressures, the amount of gas adsorbed is proportional to gas pressure.

When the gas pressure is high, $\frac{p}{y_m}$ is much larger than $1/y_m k$. Hence Eq. 2.8 could be written as,

$$\frac{p}{y} = \frac{p}{y_m}$$

or $y = y_m$ which means that at high pressures, the amount of gas adsorbed is sufficient to form a unimolecular layer. A way to verify Langmuir adsorption isotherm (Eq. 2.8) is to plot p/y against p . A straight line must be obtained. ((Fig. 2.2))




Fig. 2.2. Graphical representation of Langmuir adsorption isotherm.

It is observed that straight line plots are obtained when the surfaces are smooth and nonporous and when the pressures are not too high. Under these conditions, Eq. 2.8 is obeyed. Deviations from Langmuir adsorption isotherm are seen if;

- Surface is porous (i.e. a good adsorbent) and
- Pressure is very high.

Under these conditions, gas molecules give rise to multilayer adsorption on the solid surface which accounts for derivations from Eq. 21.8. For explaining multilayer adsorption, **Brunauer, Emmet and Teller** have proposed a model which is known as **BET** isotherm which we are not going to discuss in detail.

SELF ASSESSMENT EXERCISE 2

Starting from Eq. 2.6, derive the following equation:

$$P = \frac{0}{K(1 - 0)}$$

.....

4.4 Physisorption And Chemisorption

Depending on the type of interaction the adsorbate and the adsorbent, adsorption is of two-type-physorption (physical adsorption) and chemisorption (chemical adsorption).

Note

Both physisorption and chemisorption are generally exothermic process. That is why we talk about enthalpy decrease while comparing energies of physisorption and chemisorption.

4.41 Physisorption

If the adsorbate molecules are weakly bound to the adsorbent, it is classified as physisorption. It is also known as van der Waals adsorption since the forces involved are of van der Waals type and are of the same magnitude as are involved in the liquefaction of gases. The enthalpy decrease ($-DH$) associated with physisorption is much low (less than 40 kJ mol^{-1}) and is of the same order as the enthalpy of condensation of the adsorbate. Increase of temperature is not favourable to physisorption. Adsorption of gases by charcoal is an example of physisorption. Physisorption is generally independent of the chemical nature of the adsorbent. All gases exhibit van der Waals adsorption.

Note

The unsatisfied valencies of the surface may cause breakage of the bonds in the chemisorbed molecules. The fragments that result in the process are responsible for the increased chemical activity.

4.42 Chemisorption

If the adsorbed molecules react chemically with the surface, we call it chemisorption. The enthalpy decrease associated with chemisorption is much high (between 40 kJ mol^{-1} and 400 kJ mol^{-1}) and is of the order of bond enthalpies. We shall see in Sec. 2.8 as to how the type of bonding caused by chemisorption between the adsorbent and the adsorbate determines the reactivity pattern. Many chemisorption processes involve activation energy as in a chemical reaction. In such cases, the rates of chemisorption and desorption increase with temperature in contrast with the rate of physisorption.

Note

High bond enthalpy of nitrogen (945 kJ mol^{-1}) is mainly responsible for its low reactivity.

Let us see how the type of adsorption of nitrogen on iron surface varies with temperature. The studies on the adsorption of nitrogen on iron surface indicate that at about 770 K (the temperature chosen for Haber process), nitrogen is chemisorbed on the iron surface. Chemisorption results in large release of energy. The optimum temperature chosen for the reaction and the energy released during chemisorption are helpful in overcoming the large bond enthalpy of nitrogen. Hence, when nitrogen is chemisorbed at 770 K present as nitrogen atoms but not as molecules.

Note

One of the methods followed in scientific reasoning is to arrive at the same conclusion through more than one method. Take for instance manufacture of ammonia. Based on Le Chatelier principle, we stated in CHM 103 that high pressure ($200\text{-}300 \text{ atm}$) and optimum temperature ($670\text{-}870 \text{ K}$) are needed for a

good yield of ammonia. These conclusions could be reached from surface studies also. In this section, we have explained that around 770K, nitrogen is chemisorbed on iron to a large extent and, this facilitates formation of ammonia. As an explanation for the need of high pressure for this reaction, answer SAQ3

At temperatures less than 770K, there is not much of chemisorption of nitrogen on iron surface. At room temperature, iron does not adsorb nitrogen at all. But as temperature is lowered and brought near 80K, the boiling point of liquid nitrogen, iron adsorbs nitrogen gas physically as N₂ molecules! In short, near 770K, nitrogen is chemisorbed by iron as nitrogen atoms and near 80K, it is physisorbed as nitrogen molecules.

The dissociation of nitrogen molecule on iron surface at 770 K could facilitate its further reaction such as the formation of ammonia through Haber process. Although the mechanism of iron catalysis in Haber process is not completely understood, the chemisorption of nitrogen on iron certainly plays a role in it.

SELF ASSESSMENT EXERCISE 3

Assuming that chemisorption of nitrogen on iron at 770K follows Langmuir isotherm, could you justify the use of high pressure in Haber process for the manufacture of ammonia?

.....
.....
.....

4.5 Modern Methods of Surface Studies

The composition of the adsorbent surface, the nature of binding between the adsorbent and the adsorbate and the extent of surface coverage could be studied by using methods such as X-ray or UV photoelectron spectroscopy. Auger spectroscopy and low energy electron diffraction (LEED). Of these methods, we shall explain the principle of X-ray and UV photoelectron spectroscopy only.

Note

Auger effect is the emission of a second electron after high energy radiation has expelled on electron. Auger effect is the basis of Auger Spectroscopy and is much used in microelectronics industry.

Low energy electron diffraction is the diffraction caused by atoms on the surface by using low energy electrons. The LEED pattern depicts the two dimensional structure of a surface. Low energy electrons are used to ensure diffraction by atoms on the surface only, but not by atoms in the bulk.

In Unit 1 of Atoms and Molecules course, we have explained the photoelectric effect, according to which photoelectrons could be ejected by irradiating a metal surface with UV rays. The minimum energy that UV rays must possess for

photoelectron emission corresponds to the ionisation energy of the valence electrons. If we are interested in the emission of inner electrons, we must use X-ray or UV photoelectron spectroscopy (X.p.e.s or UV p.e.s). Since such photoelectron spectroscopy studies are useful in obtaining the finger print of the materials present in a surface of a material, these methods are known as **electron spectroscopy for chemical analysis** (ESCA). It is possible to identify the elements present in a given surface using X.p.e.s since each element has characteristic inner shell ionisation energies. The surface study using ESCA is made possible by the fact that the ejected electrons cannot escape except from within a few nanometers from the surface. The nature of chemisorption between a catalyst surface and the reactant molecules could be established by ESCA studies.

SELF ASSESSMENT EXERCISE 4

What is the basis principle for ESCA studies of the surface?

.....
.....
.....

4.6 Types of Catalysts

We have seen in Unit 18, that the rate of a chemical reaction can be increased by raising the temperature. This increases the fraction of molecules having energies in excess of some threshold energy (nearly equal to activation energy). Another way to increase the rate of chemical reaction is to find an alternate path for a chemical reaction that has a lower activation energy. A catalyst produces this alternate path for a chemical reaction. The sole function of the catalyst is to lower the activation energy of a reaction. Thus a small amount of manganese dioxide increases the rate of decomposition of KClO_3 ; the decomposition of nitrous oxide is accelerated by iodine; in presence of Ni, unsaturated hydrocarbons can be hydrogenated to saturated hydrocarbons. The amount of a catalyst remains unchanged at the end of a reaction and, may be used again and again. **A substance which can influence the rate of a chemical reaction but itself remains unchanged chemically is called a catalyst.** A catalyst cannot start a chemical reaction that could not take place in its absence. A catalyst does not alter the position of equilibrium; in other words, it cannot change the relative amounts of the reactants and products at the equilibrium. Consequently a catalyst must accelerate equally both the forward and the reverse reactions. A catalyst is highly specific in its action, e.g., MnO_2 can catalyse the decomposition of KClO_3 but not that of KNO_3 . In certain reactions, one of the products could catalyse the reaction. For instance, in the oxidation of oxalic acid by acidified KMnO_4 , Mn^{2+} ions formed during the reaction, increase the rate of reaction. This type of phenomenon is called **auto-catalysis**.

Catalysis may be of homogenous or heterogeneous type. In homogenous catalysis, the catalyst forms a single phase with the reactants and products, whereas in heterogeneous catalysis, it constitutes a separate phase.

There is another type of catalysis, known as enzyme catalysis. Enzymes have high relative molecular masses and are protein molecules. The enzymes catalyse a variety of chemical reactions in living organisms. The enzyme reaction medium is colloidal in nature and strictly speaking, enzyme catalysis does not fall under homogenous or heterogeneous catalysis. The enzymes are specific in catalyzing only a particular set of reactions. Enzyme activity depends on pH of the medium.

Examples for all the three types of catalysis are provided in Table 2.2

Table 2,2; Three Types of Catalysis

Types	Illustrative reaction	Catalyst
Homogeneous catalysis	1) $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{SO}_3(\text{g})$	NO (g)
	2) $\text{CH}_3\text{COOC}_2\text{H}_5(\text{l}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{CH}_3\text{COOH}(\text{l}) + \text{C}_2\text{H}_5\text{OH}(\text{l})$	$\text{H}_3\text{O}^+(\text{aq})$
Heterogeneous catalysis	1) $\text{HCOOH}(\text{g}) \rightarrow \text{H}_2\text{O}(\text{g}) + \text{CO}(\text{g})$	$\text{Al}_2\text{O}_3(\text{s})$
	2) $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{SO}_3(\text{g})$	Pt (s)
Enzyme catalysis	1) $\text{NH}_2\text{CONH}_2 + \text{H}_2\text{O} \rightarrow 2\text{NH}_3 + \text{CO}_2$	Urease
	2) $\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow 2\text{C}_2\text{H}_5\text{OH} + 2\text{CO}_2$	Zymase

- Note that in enzyme catalysis, we have not specified the states of the substances.

In the next section, we shall discuss the theories of catalysis.

4.7 Mechanisms of Catalytic Reactions

In homogenous or enzyme catalysis, a reaction intermediate is formed between the reactant and the catalyst or the enzyme. The intermediate compound then decomposes to give the product. The reaction sequence can be represented as follows:



Where S and P are the reactant and the product and C is the catalyst or the enzyme; SC is the intermediate compound. The role of the catalyst or the enzyme is to lower the activation energies of the forward and reverse reactions. In Fig. 21.3, E_c is the

activation energy for the conversion of a reactant to a product in the presence of a catalyst and E_{uc} is the activation energy for the same reaction in the absence of catalyst. You can see that $E_c < E_{uc}$. The same is true of the reverse reaction also.

It is interesting to note that many of the biological reactions are catalysed by enzymes. This is facilitated by the fact that the enzyme catalysed reactions have much lower activation energies than systems containing chemical catalyst. Look at Table 2.3 which indicates the activation energies for the decomposition of hydrogen peroxide under different conditions and make your own judgment!




Figure 2.1: Potential energy diagram for catalysed and uncatalysed reactions.


Table 2.3: Activation Energies for the Decomposition of Hydrogen Peroxide Solution

Catalyst	Activation energy/kJ mol ⁻¹	Relative rate of reaction
None	75.3	1
I ⁻ (aq) (homogenous)	56.5	2.0 x 10 ³
Pt (s) (heterogeneous)	49.0	4.1 x 10 ⁴
Catalase (enzyme)	8	6.3 x 10 ¹¹


Note

You can have an idea about the importance of enzyme reactions from the fact that ammonia produced from nitrogen by nitrogenase enzyme is ten times more than that produced by Haber process. Further the enzyme gives good yields of ammonia at room temperature and pressure. Compare this with the experimental conditions needed for Haber process (200-300 atm pressure and 670-870 K temperature).

In heterogeneous catalysis also, the role of the catalyst surface is to bring down the activation energies of the reactions. This happens due to chemisorption which is similar to intermediate compound formation in homogenous catalysis. The ability of a surface to have chemisorption of the reactant molecules depends on the chemical nature of surface. ESCA studies (Sec. 2.6) are helpful in deciding the nature of chemisorption between the surface and the reactant molecules. The difference in the nature of chemisorption could lead to different products even from the same reactant. For example, ethyl alcohol is dehydrogenated on Ni, Pd or Pt catalysts to give acetaldehyde.



On the other hand, ethyl alcohol undergoes dehydration reaction on alumina.



On Ni, Pd or Pt surfaces, the linkage of ethyl alcohol is through two hydrogen atoms (Fig. 2.4). The strong affinity between Ni and hydrogen accounts for the removal of two hydrogen atoms from ethyl alcohol.




Fig. 2.4: Dehydrogenation process on Ni

On the other hand, alumina acts differently due to its different structure. Alumina has both oxide groups and hydroxyl groups. The linkage of ethyl alcohol to alumina is through hydrogen and oxygen atoms as shown in Fig. 2.5. The removal of hydrogen and hydroxyl groups from adjacent carbon atoms leads to the dehydration reaction.




Fig. 2.5: Dehydration process on alumina

Certain substances improve the activity of a catalyst. Such substances are called **promoters**. These substances may not themselves be effective catalysts. A promoter may increase the number of active sites on a catalyst surface. In the light of this discussion, let us examine the catalysis in Haber process of manufacture of ammonia. A mixture of iron oxide, potassium oxide and aluminium oxide facilitates this reaction. The hydrogen atmosphere reduces iron oxide into porous iron which has large surface area that acts as the catalyst. The mixture of potassium oxide and aluminium oxide acts as promoters.

Let us now see a few applications of catalysts in chemical industries.

- In the preparation of edible fats from vegetable and animal fats, controlled partial hydrogenation with a catalyst such as nickel helps in removing some of the double bonds. In the absence of hydrogenation, these double bonds could be oxidized by air which impart the oil on rancid odour on storage.
- Careful studies of the catalytic surface have been helpful in preferential formation of a product starting from a reactant. Thus it is possible to prepare various oxidation products of ethylene such as ethanol, acetaldehyde, vinyl chloride or vinyl acetate by proper choice of catalysts and reaction conditions.
- You may be aware that ‘cracking’ is the process of producing small organic molecules by the breaking of long-chain hydrocarbon molecules. Usually, silica-alumina catalysts are used for this purpose. Cracking is required to produce branched chain isometric hydrocarbons which have more fuel efficiency in automobile engines.

SELF ASSESSMENT EXERCISE 5

At 310 K (blood temperature), sucrose could be hydrolysed using an enzyme, saccharase or a mineral acid. The reaction rate for one reaction is 10^{12} times higher than the other. Identify the faster reaction.

.....
.....
.....

SELF ASSESSMENT EXERCISE 6

In Unit 18, it has been explained that the rate of a reaction is proportional to $e^{-E/RT}$. The activation energy of the reaction $H_2(g) + I_2(g) \rightarrow 2HI(g)$ is reduced from 184 kJ mol^{-1} to 59 kJ mol^{-1} in the presence of platinum catalyst. By what factor will the reaction rate be increased by platinum at 600K? Assume that the frequency factor remains unchanged.

.....
.....

4.8 Inhibition and Poisoning

We studied that the reactants are to be adsorbed on the surface for the chemical reaction to be influenced by the surface. For the enhancement of the reaction rate, the reactants must be adsorbed more or less to same extent. If one of the reactants is more strongly adsorbed than the other or if a product is adsorbed to a greater extent than the reactants, then the active centres on the catalyst surface would not be available completely for the reaction and the reaction rate decreases. Such a condition is called inhibition of the catalyst. One of the reactants or the products which gets strongly adsorbed and thereby decreases the reaction rate is called the inhibitor. For example, in the decomposition of ammonia on platinum surface, hydrogen (a product) is strongly adsorbed and inhibits the reaction.

It is possible that the reaction could be inhibited by a foreign molecule that does not take part in a reaction. This type of inhibition is called catalytic poisoning. It is observed that even small amount of the catalytic poison could be effective in stopping a reaction. This phenomenon could be explained by the fact that the active centres constitute only a small fraction of the total surface sites on a catalyst and, the meager amount of poison could occupy these positions. This prevents the occupation by reactant molecules. For example, in the contact process of sulphuric acid manufacture, even a small amount of arsenic impurity poisons the platinised asbestos catalyst and the reaction almost stops.

SELF ASSESSMENT EXERCISE 7

State the conditions under which a catalyst loses its influence over a reaction?

.....
.....
.....

4.9 SUMMARY

In this unit, we discussed the role of surface in chemical reactions. Hence we have

- defined a surface active agent,
- discussed the significance of adsorption isotherms,
- explained physisorption and chemisorption,
- outlined the modern methods of surface studies,
- described the types of catalyst and indicated the mechanisms of their activity, and
- discussed the inhibition and poisoning of catalyst.

4.10 TUTOR MARKED ASSIGNMENT

- 1) The activation energies required for the uncatalysed decomposition of ammonia and for its decomposition in presence of tungsten are 350 and 162 kJ mol⁻¹ respectively. Explain the reason for this difference in activation energy.
- 2) The adsorption of a gas follows Langmuir isotherm with $K = 8.50 \times 10^{-4} \text{ Pa}^{-1}$ at 298 K. Find the pressure at which surface coverage is (a) 15.0% (b) 95%. Comment on your results.
- 3) The heterogeneous catalyst is often taken in the form of a finely divided powder rather than as a smooth surface. Explain the reason.
- 4) Based on adsorption studies, justify the temperature choice for Haber process.
- 5) Copper catalyses the formation of steam from hydrogen and oxygen through the formation of CuO as an intermediate. Suggest a reaction sequence for this condition.
- 6) Define the terms; (i) adsorption (ii) adsorbate (iii) adsorbent
- 7) A particular mass of charcoal adsorbs a larger volume of ammonia than of hydrogen at a given temperature. Explain.
- 8) Based on energy released during adsorption, how can you distinguish between physisorption and chemisorption?

4.11 ANSWERS TO SELF-ASSESSMENT QUESTION

- 1) The surface tension of a solution must decrease when a surface active agent is added so that its concentration is more in the surface than in the bulk.

- 2) According to Eq. 21.6,
$$0 = \frac{K_p}{1 + K_p}$$

$$0(1 + K_p) = K_p$$

$$K_p(0 - 1) = -0$$

$$K_p(1 - 0) = 0$$

$$\text{Hence, } p = \frac{0}{K(1 - 0)}$$

- 3) As per Eq. 21.6, at sufficiently high pressure $\theta = 1$. Thus at high pressure, we can expect large coverage of iron surface by nitrogen which could lead to its dissociation and also to its reaction with hydrogen to form ammonia on a large scale.
- 4) Even though X-rays may penetrate into the bulk sample, the ejected electrons cannot escape except from within a few nanometers surface. Hence this technique is mainly limited to the study of surface layers.
- 5) Enzyme catalysed reaction needs lower activation energy than the mineral acid catalysed reaction. Hence, enzyme catalysed reaction must be faster than the other.
- 6) Let r_1 and r_2 be the reaction rates in the absence of platinum and in the presence of platinum, respectively. Let E_1 and E_2 be the corresponding activation energies. Assuming frequency factors to be equal.

$$\frac{r_1}{r_2} = \frac{e^{-E_1/RT}}{e^{-E_2/RT}} = e^{(E_2 - E_1)/RT}$$

$$\text{Taking natural logarithms, } \ln \frac{r_1}{r_2} = \frac{E_2 - E_1}{RT}$$

$$\text{Or } \ln \frac{r_2}{r_1} = \frac{E_1 - E_2}{RT}$$

$$2.303 \log \frac{r_2}{r_1} = \frac{E_1 - E_2}{RT}$$

$$\text{Log } \frac{r_2}{r_1} = \frac{E_1 - E_2}{2.303RT}$$

$$\frac{r_2}{r_1} = \text{Antilog of } \frac{E_1 - E_2}{2.303RT}$$

$$= \text{Antilog of } \frac{125 \times 103 \text{ Jmol}^{-1}}{2.303 \times 8.314 \times 600 \text{ Jmol}^{-1}}$$

$$\frac{r_2}{r_1} = 7.6 \times 10^{10}$$

Hence the reaction rate is increased by 7.6×10^{10} times by platinum.

- 7) When a catalyst surface is strongly adsorbed by reactant or product or foreign gas molecules, the catalyst cannot influence the reaction rate.

UNIT 5 **PHOTOCHEMISTRY**

CONTENTS

- 5.0 Introduction
 - Objectives
- 5.1 Laws of Photochemistry
- 5.11 Grotthuss-Draper Law
- 5.12 Stark-Einstein Law of Photochemical Equivalence
- 5.2 Experimental Determination of Quantum Efficiency
 - 5.21 Thermopile
 - 5.22 Photoelectric Cells
 - 5.23 Chemical Actinometers
 - 5.24 The Quantum Efficiencies of Some Reactions
- 5.3 Photochemical Dissociation
- 5.4 Some Photochemical Reactions
 - 5.41 Photochemical Decomposition of Hydrogen Iodide
- 5.5 Photophysical Processes
- 5.6 Photosensitisation
 - 5.61 Photosensitised Chemical Reactions
 - 5.62 Sensitised Fluorescence
- 5.7 Application of Photochemistry
 - 5.71 Photogalvanic Cell
 - 5.72 Charging in presence of light
 - 5.73 Discharging in the absence of light
- 5.8 Chemiluminescence
- 5.9 Conclusion
- 5.10 Summary
- 5.11 Tutor Marked Assignment
- 5.12 References/Further Readings

5.0 INTRODUCTION

In CHM 301 we stated that free energy must decrease for a physical or a chemical process to be feasible. This is applicable to thermal reactions (also known as dark reactions). The conversion of carbon dioxide and water into starch is a reaction in which free energy increases. Under laboratory conditions, this reaction is not possible. But plants carry out starch synthesis using sunlight. Ozonisation of oxygen and decomposition of ammonia are also nonspontaneous reactions under laboratory conditions; but these could take place by using light of proper energy. Such possibilities of new reactions and new synthetic methods encouraged the scientists to probe into light-initiated reactions.

In photochemistry, we study the absorption and emission of light by matter. It consists of the study of various photophysical processes and photochemical

reactions. Two important photophysical processes are fluorescence and phosphorescence. During fluorescence, light emission takes place in the presence of exciting radiation; but the light emission stops, once the exciting radiation is removed. In contrast to this, during phosphorescence, light emission takes place even after the removal of the exciting radiation.

In photochemical reactions, the substances acquire the necessary activation energy through light absorption. Again this is in contrast to the thermal reactions in which the reactants acquire their activation energy through collisions between molecules.

In this unit, we shall discuss the laws of photochemistry. We shall then describe some photochemical reactions and photophysical processes. We shall also explain some photosensitised reactions. Finally we shall discuss the applications of photochemical studies.

OBJECTIVES

After studying this unit, you should be able to;

- explain the mathematical form of Beer-Lambert law,
- state Grotthuss-Draper law and Stark Einstein law,
- calculate the quantum yield of a reaction using the given experimental data,
- explain the reason for the higher energy requirement for the photochemical dissociation of a molecule as compared to its thermal dissociation.
- derive rate expressions for simple photochemical reactions,
- explain some photophysical properties such as fluorescence and phosphorescence,
- list some photosensitised reactions,
- state the applications of photochemical studies, and
- define chemiluminescence..

5.1 Laws Of Photochemistry

Grotthuss-Draper law and Stark-Einstein law are the two laws concerning the interaction between matter and light. Before discussing these two laws, let us state the mathematical form of Beer-Lambert law discussed in Unit 8 of Atoms and Molecules course.

Beer-Lambert law is useful in calculating the concentration of a solution on the basis of its light absorption. This law relates the intensity of the transmitted

monochromatic light to the concentration of the solution and the thickness of the cell in which the solution is kept.

Mathematical forms of Beer-Lambert law is given below:

$$\log \frac{I_0}{I} = A = \epsilon c l \quad \dots\dots 3.1$$

Where I_0 = Intensity of the incident radiation

I = Intensity of the transmitted radiation

$A = \log \frac{I_0}{I}$ = Absorbance or optical density of the solution

c = Concentration of the solution expressed in mol m^{-3} units

l = thickness of the cell

ϵ = Molar extinction coefficient; expressed in $\text{m}^2 \text{mol}^{-1}$

The molar extinction coefficient of a substance can be determined using a colorimeter or a spectrophotometer as follows. The absorbances of a solution are measured at different known concentrations using a cell of known thickness (l). The plot of A against c gives a straight line (Fig. 1.1) and its slope is equal to ϵl .




Fig.1.1 A vs c plot.

Hence, $\epsilon = e \frac{\text{slope}}{l}$

Since l is known, ϵ can be calculated. Using this ϵ value, the concentration of a solution can be determined by measuring its absorbance.

Colorimeters and spectrophotometers are commercially available for the measurement of adsorbances. In colorimeters, colour filters are used to get the incident radiation in a particular wavelength range. Spectrophotometers have arrangements for obtaining nearly monochromatic incident radiation. Both colorimeters and spectrophotometers have devices for converting the light transmitted into suitable signal through detector devices (see Sec. 3.3). The signal generated from transmitted light is directly read as absorbance values.

Study the following example carefully. The calculation shown below is used in chemical actinometers (Example 3 in Sec. 3.3) for estimating the intensity of the light absorbed during a photochemical reaction. The example illustrates the use of Eq. 1.1 in calculating the concentration of a solution.

Example 1

A solution of the red complex formed by Fe^{2+} ion with 1,10-phenanthroline is taken in a cell of the thickness 1.00cm. If ϵ for the complex is $1.11 \times 10^3 \text{ m}^2 \text{ mol}^{-1}$ and the absorbance of the solution is 0.391, calculate the concentration of the complex in mol m^{-3} units.

Solution


$$A = 0.391; l = 1.00 \text{ cm} = 1.00 \times 10^{-2} \text{ m};$$

$$\epsilon = 1.11 \times 10^3 \text{ m}^2 \text{ mol}^{-1}$$

$$\text{Rearranging Eq. 3.1, } c = \frac{A}{\epsilon l} = \frac{0.391}{1.11 \times 10^3 \times 1.00 \times 10^{-2}} \text{ mol m}^{-3}$$

$$= 0.0352 \text{ mol m}^{-3}$$

Note



1,10 – Phenanthroline (Phen)

We shall now state the two laws of photochemistry.

5.11 Grotthuss-Draper Law

According to this law, only the light that is absorbed by a molecule can produce a photochemical change in it. This means that it not sufficient to pass through a substance to bring about a chemical reaction; but the light must be absorbed by it.

Stark-Einstein law of photochemical equivalence provides a quantum mechanical form to Grotthuss-Draper law.

5.12 Stark-Einstein Law of Photochemical Equivalence

Stark-Einstein law of photochemical equivalence can be stated as follows:

Each molecule taking part in a photochemical reaction absorbs one quantum of radiation which causes the reaction.

Note

Stark-Einstein law is applicable only if the intensity of light is not very high.

This law is applicable to the primary act of excitation of a molecule by light adsorption. This law helps in calculating the quantum efficiency (F) which is a measure of the efficiency of the use of light in a photochemical reaction.

The quantum efficiencies for the formation of a product and for the disappearance of the reactant are defined below:

The quantum efficiency
For the formation of Product (fR) = $\frac{\text{Number of molecules of product formed in 1 second}}{\text{Number of quanta absorbed In 1 second}}$ 3.2

$$= \frac{dN_p / dt}{I_a}$$

In this equation, dN_p/dt denotes the rate of formation of the product. The units for $\frac{dN_p}{dt}$ are molecules per second. I_a refers to the number of photons absorbed per second. Similarly we can define the quantum efficiency for the disappearance of a reactant.

The quantum efficiency
For the disappearance of the reactant (fR) = $\frac{\text{Number of molecules of reactant consumed in 1 second}}{\text{Number of quanta absorbed In 1 second}}$ 3.3

$$= \frac{-dN_r / dt}{I_a}$$

Where $-dN_r/dt$ is the rate of consumption of the reactant. The minus sign in $-dN_r/dt$ is due to the decrease in concentration of the reactant with time.

As per Stark – Einstein law, each reacting molecule absorbs one quantum of light.
Hence,

The energy absorbed by one
Mole of substance undergoing

$$\text{Photochemical reaction} = N_A h \nu = \frac{N_A hc}{\lambda}$$

This unit of energy is also called Einstein and, as apparent from Eq. 19.4, the value of Einstein depends on ν or λ .

Where $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

$$h = 6.626 \times 10^{-34} \text{ J s}$$

$$c = 3 \times 10^8 \text{ m s}^{-1}$$

λ = wavelength of the light in m unit

ν = frequency in s^{-1} unit.

To calculate the quantum efficiency, use the following steps:

i) Calculation of Energy Absorbed per Quantum

$$\begin{array}{l} \text{Energy per quantum} \\ \text{of radiation absorbed} \end{array} = h\nu = \frac{hc}{\lambda}$$

ii) Calculation of I_a

Usually intensity of light absorbed is given in J s^{-1} units. From this, I_a could be calculated using the following equation:

$$\begin{array}{l} \text{Intensity in terms of number of} \\ \text{photos absorbed in 1 second (1)} \end{array} = \frac{\text{Intensity in } \text{J s}^{-1} \text{ units}}{\text{Energy per quantum}}$$

Note:

Unit for energy per quantum = joule photon⁻¹

$$\text{Unit for } I = \frac{\text{joule second}^{-1}}{\text{joule photon}^{-1}}$$

$$= \text{photon second}^{-1}$$

iii) **Calculation of** $-\frac{dN_R}{dt}$ or $\frac{dN_P}{dt}$

First we have to obtain the number of moles of reactant consumed or of product formed **in 1 second** by dividing the respective amount by the time in second unit.

In order to calculate the rate of formation or rate of disappearance in molecule per second units, we have to use the following equations:

$$\frac{-dN_R}{dt} = \begin{array}{l} \text{Number of moles of reactant} \\ \text{Decomposed in 1 second} \end{array} \times N_A \quad \dots 3.5$$

$$\text{Also, } \frac{dN_P}{dt} = \begin{array}{l} \text{Number of moles of product} \\ \text{Formed in 1 second} \end{array} \times N_A \quad \dots 3.6$$

iv) **Calculation of Quantum Efficiency**

Using Eq. 3.2 or 3.3 f_p or f_R can be calculated.

Let us illustrate the calculation of quantum efficiency for a reaction.

Example 2

In Photochemical decomposition of acetone using 313 nm light, 7.57×10^{-6} mol of carbon monoxide is formed in 20 minutes. If the light absorbed corresponds to $2.41 \times 10^{-3} \text{ J s}^{-1}$, calculate the quantum efficiency for the formation of carbon monoxide.

$$\text{i) energy absorbed per quantum} = \frac{hc}{\lambda} = \frac{6.626 \times 10^{-34} \times 3 \times 10^8}{313 \times 10^{-9}} \text{ joule photon}^{-1}$$

$$= 6.35 \times 10^{-19} \text{ joule photon}^{-1}$$

$$\text{ii) } I_a = \frac{\text{Intensity in joule second}^{-1}}{6.35 \times 10^{-19} \text{ joule photon}^{-1}}$$

$$= \frac{2.14 \times 10^{-3} \text{ joule second}^{-1}}{6.35 \times 10^{-19} \text{ joule photon}^{-1}}$$

$$\text{i.e. , } I_a = 3.80 \times 10^{15} \text{ photons second}^{-1}$$

$$\begin{aligned}
 \text{iii) Amount of carbon monoxide} &= \frac{7.57 \times 10^{-6} \text{ mol}}{20 \times 60 \text{ s}} \\
 \text{formed in 1 second} & \\
 &= 6.31 \times 10^{-9} \text{ mol s}^{-1} \\
 \frac{dN_{\text{CO}}}{dt} &= 6.31 \times 10^{-9} \times 6.022 \times 10^{23} \text{ molecule second}^{-1} \\
 &= 3.80 \times 10^{15} \text{ molecule second}^{-1}
 \end{aligned}$$

$$\begin{aligned}
 \text{iv) Using Eq. 3.2 f CO} &= \frac{\text{Quantum efficiency for the}}{\text{formation of carbon monoxide}} \\
 &= \frac{dN_{\text{CO}}}{I_a} = 1
 \end{aligned}$$

In the next section, we shall discuss the experimental method of determining the quantum efficiency of a reaction. Before studying this, work out the CHM.

SELF ASSESSMENT EXERCISE 1

A radiation of 250nm incident in H1 results in the decomposition 1.85×10^{-2} mol within a particular time interval. Light energy absorbed during this time interval is 4.18 k J.

Calculate the quantum efficiency for the decomposition of H1.

.....

5.2 Experimental Determination of Quantum Efficiency

For the determination of quantum efficiency of a photochemical reaction, we must measure the following:

- the rate of decomposition of the reactants or the rate of formation of the products; this can be done using any of the methods discussed in Units 18.
- the number of quanta absorbed in the given time.

In short, we need a method for measuring the amount of light absorbed, if we can devise a method for estimating the intensity of the incident light (I_o) and the intensity of the light transmitted (I) by the reaction mixture, the intensity of the light absorbed (I_a) can be calculated using the relationship;

$$I_a = I_o - I$$

For measuring I_0 and I , same cell is used. I_0 corresponds to the measurement of intensity with empty cell (known as blank) and I , to the measurement using the cell containing the reaction mixture.

The experimental set-up used for measuring the intensity of light is shown in Fig. 3.2

As a source (A) of visible light, a high intensity tungsten lamp or quartz halogen lamp or xenon arc lamp can be used. For obtaining ultraviolet light, hydrogen or deuterium lamps are good sources. The light is first passed through a lens, B (to get a parallel beam of light), and then through a monochromator (C). The monochromator yields a narrow band of light in the desired wavelength. The monochromatic light then passes through the reaction cell (D). The light transmitted by the reaction cell reaches the detector (E).




Fig 3.2: Measurement of Intensity of light during a photochemical reaction

Note

While using ultraviolet light as a source of radiation, quartz cuvettes (cells) and lenses must be used.

Since pyrex glass absorbs light below 300nm, it can be used for radiation above 300nm only

The visible light has the wavelength region, 400-700nm. The ultraviolet light has wavelength below 400nm and above 100nm.

The detector is used for measuring the intensity of light. The quantitative measurement of light intensity based on energy conversion or on chemical reaction is called **actinometry**. Detectors such as thermopiles and the photoelectric cells function on energy conversion principle. We discuss each of them in brief below:

5.21 Thermopile

Thermopile (Fig 3.3) is a device useful for converting light into heat, and then, heat into electricity. It has a set of junctions of different metals having a blackened surface. This black coating is to ensure absorption of all radiation falling on it. The energy so absorbed increases the temperature of the metals. Thermopile converts this temperature increase into a potential difference. Light intensity is related to electricity generated by it. The detector device must be calibrated against a standard light source. Thermopile could be used as a detector for light of any wavelength.

Note:

Thermopile consists of thermocouples connected in series and it generates electricity on heating





Fig. 3.3: Thermopile: I. light radiation, II. Blackened surface exposed to radiation; III. unexposed surface at the back; IV galvanometer.

5.22 Photoelectric Cells

Photoelectric cell converts light directly into electricity. The current generated is directly proportional to light intensity. Photoelectric cell is an evacuated bulb with a photosensitive cathode. Light radiation results in the emission of electrons from the cathode which get collected at the anode. Thus the flow of current is caused. The photoelectric cell is sensitive to the wavelength of the light used and must be calibrated using a thermopile.

5.23 Chemical Actinometers

Light intensity could be estimated by a quantitative estimation of a substance formed or decomposed by light. The photochemical decomposition of $\text{K}_3\text{Fe}(\text{C}_2\text{O}_4)_3$ is a widely used chemical method for estimating intensity of light between the wavelength range 250nm and 577nm. Light within this wavelength range causes $\text{K}_3\text{Fe}(\text{C}_2\text{O}_4)_3$ dissolved in sulphuric acid to react as follows:



The concentration of Fe^{2+} is estimated colorimetrically through complex formation with 1,10-phenanthroline. (See Example 1). This complex has a deep red colour. The concentration of the complex is equal to the concentration of Fe^{2+} present before complexation. The following example is worked out to illustrate this method.

Example 3

In a potassium ferrioxalate actinometer, the quantum efficiency for Fe^{2+} production (f FS) at 480 nm wavelength is 0.95. After irradiating the potassium ferrioxalate solution for 20 minutes, it is completely transferred to a 200cm³ volumetric flask, mixed with required quantity of 1,10-phenanthroline for complex formation and made up to mark with a buffer solution. For colorimetric estimation, a sample of this complex is taken in a cell of 1.00cm thickness. The complex has an absorbance value of 0.391 and its ϵ is $1.11 \times 10^3 \text{ m}^2 \text{ mol}^{-1}$.

Calculate

- i) the concentration of complex,
- ii) the number of Fe^{2+} ions formed in the actinometer due to irradiation,
- iii) the rate of formation of Fe^{2+} ions (dN_{FS}/dt)
- iv) I_a (number of photons absorbed per second)

Assume that one Fe^{2+} ion forms one complex molecule with 1,10-phenanthroline.

Solution

- i) Note that the concentration of the complex is same as that calculated in Example using Eq. 1.1; $c = \text{concentration of complex} = 0.0352 \text{ mol m}^{-3}$

Also, concentration of Fe^{2+} ion = concentration of the complex

$$\begin{aligned}
 &= 0.0352 \text{ mol m}^{-3} \\
 \text{ii) } &\text{Number of moles of Fe}^{2+} \text{ Present in } 200\text{cm}^3 \text{ of the solution} = \frac{\text{Concentration of Fe}^{2+} \times \text{Volume of the Solution in m}^3 \text{ unit}}{\text{of Fe}^{2+}} \\
 &= 0.0352 \times 200 \times 10^{-6} \text{ mol} \\
 &= 7.04 \times 10^{-6} \text{ mol}
 \end{aligned}$$

Because $200\text{cm}^3 = 200 \times 10^{-6} \text{ m}^3$
Hence the number of moles Fe^{2+} formed in the actinometer = $7.04 \times 10^{-6} \text{ mol}$

The number of Fe^{2+} ions formed in the actinometer = $7.04 \times 10^{-6} \times 6.022 \times 10^{23}$ ions
= 4.24×10^{18} ions

$$\begin{aligned}
 \text{iii) } \frac{dN_{\text{FS}}}{dt} &= \frac{\text{Number of Fe}^{2+} \text{ ions formed}}{\text{Time in seconds}} = \frac{4.24 \times 10^{18}}{1200} \text{ ions second}^{-1} \\
 &= 3.53 \times 10^{15} \text{ ions second}^{-1}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) } \text{Using Eq. 3.2} &= \frac{dN_{\text{FS}}/dt}{l_a} \\
 \text{Given that } f_{\text{FS}} &= 0.95 \\
 \text{Hence } l_a &= \frac{dN_{\text{FS}}}{dt} \times \frac{1}{0.95} \\
 &= \frac{3.53 \times 10^{15}}{0.95} \text{ photons second}^{-1} \\
 &= 3.72 \times 10^{15} \text{ photons second}^{-1}
 \end{aligned}$$

5.24 The Quantum Efficiencies of Some Reactions


If a photochemical reaction follows Stark-Einstein law, then f must be unity. Studies on a large number of photochemical reactions indicate that while some of them follow Stark-Einstein law, many others do not. The quantum efficiencies of some photochemical reactions along with their effective wavelengths are given in Table 3.1

Table 3.1: Quantum Efficiencies and Effective Wavelengths of Some Photochemical Reactions

Reaction	Effective wavelength/nm	f
$2\text{NH}_3 \rightarrow \text{N}_2 + 3\text{H}_2$	210	0.2
$2\text{NO}_2 \rightarrow 2\text{NO} + \text{O}_2$	366	0.7
$\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$	511	0.01
$\text{H}_2\text{S} \rightarrow \text{H}_2 + \text{S}$	208	1.0
$2\text{HI} \rightarrow \text{H}_2 + \text{I}_2$	327	2.0
$3\text{O}_2 \rightarrow 2\text{O}_3$	175	3.0
$\text{H}_2 + \text{Cl}_2 \rightarrow 2\text{HCl}$	478	10^6

The photochemical reactions for which quantum efficiency is a small integer are said to follow Stark-Einstein law. Examples are the dissociation of H_2S or ozonisation of oxygen etc. Reactions like the formation of hydrogen chloride or hydrogen bromide do not follow Stark-Einstein law. The quantum efficiency for the former reaction ($\text{H}_2 + \text{Cl}_2 \rightarrow 2\text{HCl}$) is very high while it is much low for the latter ($\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$). In order to explain these discrepancies, Bodenstein pointed out that photochemical reactions involved two distinct processes:

- i) In the primary process, absorption of one quantum of light results in the excitation of one molecule or an atom (A) to form the excited species, A^{*}



- ii) In the secondary process, the excited species undergoes a chemical change



According to Bodenstein, the deviation from Stark-Einstein law are due to the secondary processes.

In the case of reactions having low quantum yields, the number of molecules decomposed by the absorption of one quantum of radiation is less than one. The probable reasons for the low quantum yields are given below:

- The excited species formed at the first step may be deactivated by collisions before the product could be formed. This process is called **quenching**. Sometimes the presence of paramagnetic substances like oxygen or nitric oxide could result in quenching leading to low quantum yield for a chemical reaction.
- One or more of the reactions in the secondary/processes may be endothermic. High energy requirement could decrease the reaction rate. In Sec. 3.5, we will see that the quantum yield for the formation of hydrogen bromide is low due to the endothermic nature of the reaction between Br and H_2 (see Example 5 in Sec. 3.5).

- The excited species may recombine to give the reactant molecule. In the hydrogen bromide formation, another reason for the low quantum yield is the recombination of bromine atoms which are formed in the primary step.

In the case of reactions with high quantum yields, the excited species formed in the first step could initiate a series of chain reactions. This causes a large number of molecules to react by absorption of one quantum of light, as in the case of hydrogen chloride formation (see Sec. 3.5).

The following CHM could be answered correctly, if you have understood the materials in this section.

SELF ASSESSMENT EXERCISE 2

The quantum yield for the photochemical formation of hydrogen chloride is high in the absence of oxygen but low when oxygen is present. Explain.

.....

5.3 Photochemical Dissociation

In CHM 301 it has been stated that the bond enthalpy gives an estimate of the average energy required to break a particular bond. These bond enthalpies are derived from thermochemical calculations (see unit 1 of CHM 301). It is generally seen that the energy required for the dissociation of a particular bond by light absorption is much greater than is bond enthalpy value. In Table 3.2, we illustrate this for some diatomic molecules.

Table 3.2: Comparison of Photochemical Dissociation Energies and Bond Enthalpies

Dissociating molecules	λ /nm	(Photochemical dissociation''energy) kJmol^{-1}	Bond enthalpy'' kJ mol^{-1}
H ₂	84.5	1420	436
Cl ₂	478	250	242
Br ₂	511	235	193
I ₂	327	367	299
O ₂	176	682	497

Note


Bond enthalpy of a diatomic molecule indicates the energy requirement for its thermal dissociation.

* λ value corresponds to maximum wavelength of light required for decompositions by direct irradiation.

** Photochemical dissociation energy for one mole is calculated by substituting for l in Eq. 3.4

The reasons for the higher photochemical dissociation energies as compared to bond enthalpies are given below:

- For dissociating a molecule through light absorption, there must be an upper electronic state with appropriate energy levels. There is no such restriction for thermal decomposition.
- During photochemical dissociation, the product species could be an excited state and/or in ground state. For example, decomposition of bromine by absorption of light of wavelength 511 nm can be shown as below:



The atom with asterisk sign indicates excited state. Thus, photochemical decomposition of bromine needs 235 kJ mol^{-1} . But thermal decomposition of bromine needs 193 kJ mol^{-1} only, since both the bromine atoms are formed in ground state.

The photochemical dissociation of molecules is also known as photolysis. The photolysis can be understood using potential energy diagrams.

In Fig. 3.4, the ground state and the excited state are represented using potential energy diagrams I and II, respectively. The quantized vibrational sublevels in each state are shown by horizontal lines such as AB, EF etc.




Fig. 3.4: Electronic excitation; E (potential energy) vs. r (internuclear distance)


When a molecule is excited from zero vibrational level AB of ground electronic state of any of the vibrational levels **below GH** in the upper electronic state, the resultant electronic spectrum shows an absorption band with vibration – rotation fine structure. The fine structure is due to numerous transitions (such as III, IV etc.) possible from the zero vibrational level (AB) in the lower electronic state to any of the quantized vibrational levels in the upper electronic state. This can further be understood from the fact that each vibrational level has its own rotational sublevels.

When a molecule absorbs sufficient energy such that it is transferred from the ground state to or above GH in the upper electronic state, then the molecule undergoes photochemical dissociation. The spectrum shows a continuum (lack for discrete lines), once the molecule dissociates. The difference in energy between the levels AB and GH (E_p) is the **photochemical dissociation energy**. The **thermal dissociation energy** (E_T) is equal to the bond enthalpy in the case of diatomic molecules and it is the energy difference between the lowest and uppermost vibrational levels (AB and CD) in the ground state. Note that $E_p > E_T$.

We have disused photolysis in detail so far because it is initial reaction in many photochemical reactions. The excited atom or radical formed due to photolysis of a molecule often starts a chain reaction. We shall use this principle in the next section in the study of some photochemical reactions. Before studying that, answer the following SSQ which will help you in understanding Example 4.

SELF ASSESSMENT EXERCISES 3

In the photochemical dissociation of HI, the first step is given below:



Note

The excess energy ($\Delta E_p - \Delta E_T$) that a photochemical decomposition demands as compared to thermal decomposition is given to one of the atoms formed. This atom is said to be in the excited state.

Assume that H atom formed is in excited state while I atom is in ground state. Calculate the excess energy that the excited hydrogen atom carries as (use Table 19.2) compared to a ground state hydrogen atom.

.....
.....
.....

5.4 Some Photochemical Reactions


In this section, we discuss the mechanism of some photochemical reactions and then explain flash photolysis. For the first two reactions given below, we derive rate equations also. You go through these derivations carefully. These two examples could give you an ideal as to how the photochemical rate expressions are written.

The first step in both the examples is photolysis. The rate of the photolysis step is expressed as I_a which is the rate of absorption of light (number of quanta absorbed per second). The initial photolysis is followed by thermal (or dark) reactions for which kinetic expressions as similar to those discussed in Unit 18.

Example 4

5.41 Photochemical Decomposition of Hydrogen Iodide

Let us derive expressions useful in calculating the rate of decomposition of HI and the quantum efficiency for this reaction. HI undergoes photochemical decomposition below 327 nm. The mechanism is given below:



Such steps are written based on energy considerations, if you have answered CHM in the last section correctly, you could follow why in the second step H atom attacks HI whereas I does not.

Note

While deriving rate expressions, we try to eliminate terms contain active species using steady state principle.

HI is consumed in two ways as per Eqs. 3.7 and 3.8. The rate of disappearance of HI can be written as follows:

$$-\frac{d[\text{HI}]}{dt} = I_a + k_2 [\text{H}] [\text{HI}] \quad \dots 3.10$$

As per steady state approximation discussed in CHE 311 the concentration of the active species H is constants. In other words, its concentration does not vary with time.

$$\text{i.e., } \frac{d[\text{H}]}{dt} = 0 = I_a - k_2 [\text{H}] [\text{HI}] \quad \dots 3.11$$

Note that H is formed as per Eq. 3.7 and is used as per Eq. 3.8

$$\text{Or } k_2 [\text{H}] [\text{HI}] = I_a \quad \dots 3.12$$

Using Eqs. 3.10 and 3.12, we get,


$$\frac{-d[\text{HI}]}{dt} = 2 I_a \quad \dots 3.13$$

Experimentally it has been observed that the quantum efficiency for HI decomposition is 2 (Have you worked out CHM 1?). It is worth nothing that the rate of decomposition of HI depends on the intensity of the absorbed light as per Eq. 3.13.

Example 5

Photochemical Reaction between H₂ and Br₂

Hydrogen and bromine combine at wavelengths below 511 nm to give HBr as per the mechanism given below. Let us calculate the rate of formation of HBr and also the quantum efficiency for this reaction.



The rate of formation of HBr can be written as,

$$\frac{d[\text{HBr}]}{dt} = k_2 [\text{Br}] [\text{H}_2] + k_3 [\text{H}] [\text{Br}_2] - k_4 [\text{H}] [\text{HBr}] \quad \dots 3.19$$

Note that Br and H are active species and we can apply steady state approximation for these two.

$$\frac{d[\text{H}]}{dt} = 0 = 2I_a - k_2 [\text{Br}] [\text{H}_2] + k_3 [\text{H}] [\text{Br}_2] - k_4 [\text{H}] [\text{HBr}] - 2k_5 [\text{Br}] \quad .3.20$$

The coefficient 2 in $2I_a$ and $2k_5 [\text{Br}]$ are due to the formation or disappearance of two bromine atoms in the respective steps.

$$\frac{d[\text{H}]}{dt} = 0 = k_2 [\text{Br}] [\text{H}_2] - k_3 [\text{H}] [\text{Br}_2] - k_4 [\text{H}] [\text{HBr}] \quad \dots 3.21$$

Adding Eq. 3.20 and using Eq 3.21 and rearranging we get,

$$2 k_5 [\text{Br}]^2 = 2I_a \quad \dots 3.22$$

$$\text{Or} \quad [\text{Br}] = (I_a / k_5)^{1/2} \quad \dots 3.23$$

Rearranging Eq. 3.21 and using Eq. 3.23 we get,

$$[\text{H}] = \frac{k_2 [\text{Br}] [\text{H}_2]}{k_4 [\text{HBr}] + k_3 [\text{Br}_2]} = \frac{k_2 [\text{H}_2] (I_a/k_5)^{1/2}}{k_4 [\text{HBr}] + k_3 [\text{Br}_2]} \quad \dots 3.24$$

Using Eqs. 3.19, 3.23 and 3.24

$$\frac{d[\text{HBr}]}{dt} = k_2 (I_a/k_5)^{1/2} [\text{H}_2] + (k_3 [\text{Br}_2] - k_4 [\text{HBr}]) \frac{k_2 [\text{H}_2] (I_a/k_5)^{1/2}}{k_4 [\text{HBr}] + k_3 [\text{Br}_2]} = \frac{2k_2 k_3 (I_a/k_5)^{1/2} [\text{H}_2] [\text{Br}_2]}{k_4 [\text{HBr}] + k_3 [\text{Br}_2]}$$

Dividing the numerator and the denominator of R.H.S by $k_3 [\text{Br}_2]$

$$\frac{d[\text{HBr}]}{dt} = \frac{2k_2 k_5^{-1/2} I_a [\text{H}_2]}{1 + k_4 [\text{HBr}] / k_3 [\text{Br}_2]} \quad \dots 3.35$$

$$\text{i.e.,} \quad \frac{d[\text{HBr}]}{dt} \propto I_a^{1/2} \quad \dots 3.26$$

This shows that the rate of hydrogen bromide formation is proportional to the square root of absorbed intensity. This has been proved experimentally.

In the above two cases, we derived rate expressions. Next, we shall study the reaction between H_2 and Cl_2 in a qualitative way, and then, study the principles of flash photolysis.

Hydrogen-chloride Reaction

The quantum efficiency for the photochemical combination of hydrogen and chlorine is very high. When exposed to light of wavelength 478nm, the primary process in the photochemical reactions is the decomposition of chlorine.



This is followed by the following secondary processes:



The reaction between Cl and H_2 (Eq. 3.28) is exothermic and much fast. This results in the propagation of chain reaction with high quantum efficiency. The chain-terminating step is the recombination of chloride atoms on the walls of the vessel to form chloride molecule.



Next we shall explain briefly the main features of flash photolysis. Flash photolysis is useful in detecting the short-lived intermediates in a reaction sequence. The reaction mechanisms are proposed based on flash photolysis results.

Flash photolysis


Flash photolysis was developed by Norrish and Porter in 1949. In ordinary photolysis, the steady state concentrations of the intermediates are so small that these cannot be detected by absorption spectrophotometers. In flash photolysis, a high-intensity flash of microsecond duration is used for photolysis the substance and the products are identified using absorption spectrophotometers. The flash duration must match the decay rate of the intermediates. Flash lamps work for a time of around 15 μs . This restricts their use to the study of intermediate of life time around 100 μs . In recent years, laser flash sources have been developed. The flash duration is around 10^{-9} s.

As far as this section is concerned, make sure that you understand the derivations for calculating the rate of decomposition of hydrogen iodide and the rate of formation of hydrogen bromide. This could help you in arriving at the rate expressions for simple photochemical reactions for which reaction sequence (like

Eq. 3.14 to Eq.3.18) is known. You can build up confidence by answering the CHM given below. The aim of this CHM is to make you derive expression for the rate of formation of carbon monoxide in the photolysis of acetaldehyde. You are guided through a series of steps with helpful hints. This guidance has added to the length of the problem. Don't mind it!

SELF ASSESSMENT EXERCISE 4

Look at the reaction sequence for the photolysis of acetaldehyde:



I_a is the absorbed light intensity and it represents the rate of photochemical excitation (as per Eq.(1)).

Using the above mechanism, derive expressions for the following:

i) $\frac{d[\text{CH}_3\text{CO}]}{dt}$ ii) $\frac{d[\text{CH}_3]}{dt}$

(Hints: CH_3CO and CH_3 are active species. CH_3CO is formed as per Eq. (2) and consumed as per Eq. (3). CH_3 is formed as per Eqs. (1) and (3), and used up as per Eqs. (2) and (4). The answers for (i) and (ii) are to be arrived at using Eqs. 3.21 and 3.22 as models. Note that as per Eq. (4), two CH_3 radicals are consumed for every molecule of ethane formed).

iii) Derive the combined expression for $\frac{d[\text{CH}_3\text{CO}]}{dt} + \frac{d[\text{CH}_3]}{dt}$

(Hints: Add up the expressions you have got as answers for (i) and (ii).

Note

For a 2-election system, the first three singlet states are represented below:

iv) Find the relationship between $[\text{CH}_3]$ and I_a

(Hint: Rearrange the answer for (iii)

v) State the relationship between $\frac{d[\text{CO}]}{dt}$ and $[\text{CH}_3\text{CO}]$


(Hint: Use Eq. (3).)

vi) Find the relationship between the rate of formation of CO and the absorbed light intensity.

(Hint: First combine answers for (v) and (i). Then use the answer for (iv))


Note

For a 2-electron system, the first three singlet states are represented below:



Note that $E_{s2} > E_{s1}$ where E stands for the energy of a particular level.

Similarly, the first three triple states are represented below for a 2-electron system:



.....

5.5 Photophysical Processes

The light absorbed by a molecule is not always used up in producing a chemical reaction. The absorbed energy can be lost through various physical processes also. In this section, we examine such physical processes.

The adsorption of ultraviolet or visible light results in the increased of electronic energy from the ground to the excited stated. Usually electronic excitation is also accompanied by an increase in the rotational and vibrational energy levels. In our discussion, for convenience, we depict only transitions between electronic energy levels. In order to understand the nature of electronic transitions, it is essential to know the concept of spin multiplicity. A molecule with electrons pairs and with anti-parallel spins is said to be in single ground state (S_0). An excited molecule with **two of its electrons unpaired and, with anti-parallel spins** is said to be in the excited singlet state such as S_1, S_2, S_3, \dots etc. An excited molecule with **two of its electrons unpaired, but with parallel spins** is said to be in the excited triple state such as T_1, T_2, T_3, \dots etc.

Note

Normally excitation of a ground state molecule leads only to one of the excited singlet states. But in some specified cases, direct excitation from S_0 to a triplet level is possible. We restrict our discussion to $S_0 \rightarrow S_1, S_0 \rightarrow S_2$ etc, transitions only.

Multiplicity of state is given by the expression $2S + 1$, where S (note the italicized type) is the sum of spin values of electrons. This symbol S should not be confused with S (Roman type) for singlet state.

Imagine a molecule in the singlet state. It has two electrons with anti-parallel spins (such as $\uparrow\downarrow$). Then, sum of the electron spins

$$= S = +\frac{1}{2} - \frac{1}{2} = 0$$

Hence multiplicity

$$= 2S + 1 = (2 \times 0) + 1 = 1$$

Normally a molecule in the S_0 state on absorbing a quantum of light gets two of its paired electron unpaired and gets transferred to S_1 or S_2 or S_3, \dots etc, levels, but not to T_1 or T_2 or T_3, \dots etc levels. That is, due to excitation, spin multiplicity is not generally altered. This condition is called the selection rule for electronic transition. In other words, absorption of energy by the molecule in the ground state leads to allowed transition such as $S_0 \rightarrow S_1, S_0 \rightarrow S_2, S_0 \rightarrow S_3$ and so on. Such excitations and the subsequent energy loss while reaching the ground state are shown by Jablonski diagram (Fig. 3.5). The solid arrows pointing upwards refer to absorption of energy. The solid arrows pointing downwards refer to energy emission as light, known as radiative transition. The wavy horizontal arrows stand for transition between excited singlet and triplet state without energy loss, while wavy vertical arrows stand for transition between singlet-singlet or triplet-triplet levels with energy loss as heat (the later is not shown in Fig. 3.5). These wavy arrows stand for nonradiative transition (transitions without light emission).

Note

Thus the singlet state has two electrons with anti-parallel spins and its $(2S + 1)$ value is equal to 1. In the presence of a magnetic field, a singlet state does not split further.

A molecule in the triplet state has two electrons with parallel spins (such as $^{\circ\circ}$). The sum of electron spins.

$$= S = \frac{1}{2} + \frac{1}{2} = 1$$

Hence the multiplicity $= (2S + 1) = 3$

So, a molecule in the triplet state has two electrons with parallel spins and its $(2S + 1)$ value is equal to 3. In the presence of a magnetic field, a triplet state splits into three energy levels.

In Fig. 3.5, the excitation from the singlet ground state to the excited singlet levels.




Fig. 3.5: Jablonski diagram

S_1 , S_2 and S_3 are shown by vertical arrows marked A. The excited species at S_2 and S_3 have very short lifetimes and these species, quickly lose their energy as heat to the medium in about 10^{-11} second and reach S_1 level. Such a singlet-singlet transition is called an internal conversion (IC). The molecule at S_1 state has a life time 10^{-8} - 10^{-10} second. The system at S_1 may undergo any of the following transitions.

A = Absorption of quanta leading to excitation to S_1 , S_2 and S_3 levels. Note the antiparallel spins at various S levels

IC = Internal conversion from S_3 to S_2 , from S_2 to S_1 and from S_1 to S_0


F = Fluorescence; a transition from $S_1 \rightarrow S_0$ with light emission.

ISC = Intersystem crossing; $S_1 \rightarrow T_1$ transition. Note the parallel spins at various T levels

**P = Phosphorescence; a transition from $T_1 \rightarrow S_0$ with light emission
For clarity, transition to and from T_2 and T_3 are not shown.**

Note

The name “fluorescence” is derived from the name of the mineral, “fluorite”, which emits visible light on exposing to ultraviolet radiation.



Benzophenone

i) Fluorescence

The excited molecule could undergo the transition, $S_1 \rightarrow S_0$, with emission of light. This phenomenon is called fluorescence (F). Since $S_1 \rightarrow S_0$ transition is allowed by selection rule, it is very fast. In other words, substances fluoresce in the presence of the exciting radiation. Once the exciting radiation is stopped, the fluorescence stops.

ii) Internal Conversion

The excess energy may be lost as heat while $S_1 \rightarrow S_0$ transition takes place which is again a case of internal conversion and a radiationless transition.

iii) Intersystem Crossing

The excited molecule could cross over to the first triplet state through $S_1 \rightarrow T_1$ transition. Such a transition involves spin inversion. For such intersystem crossing (ISC) to be efficient, the energy gap between S_1 and T_1 levels must be low.

Ketones have very low energy gap between S_1 and T_1 levels and have high efficiency for intersystem crossing. Thus benzophenone has 100% efficiency for intersystem crossing. Compared to ketones, aromatic hydrocarbons are less efficient in intersystem crossing and olefins are still less efficient. Let us now study two of the processes by which the molecule in the triplet state could reach the ground state.

1) Phosphorescence

An interesting physical process by which an excited species at T_1 level may undergo transition to S_0 level is by emitting light; $T_1 \rightarrow S_0$ transition with emission of light is called **phosphorescence (P)**. This is a process with a spin change and is a forbidden transition. Hence, in contrast to fluorescence, light emission during phosphorescence is slow and it lasts even after the removal of exciting radiation.

2) Energy Transfer

Another process by which a molecule in the triplet state (called a donor molecule) may lose its excess energy is by energy transfer to an acceptor molecule. This is an instance of **sensitisation** which we shall discuss in Sec. 3.7.

In this section, we have dealt with some of the important physical processes only. Let us now see some of the applications of the study of the physical processes. Study of fluorescent behaviours of substances has led to the development of fluorescence spectroscopy. Using spectrofluorometers, it is possible to identify several fluorescing substances present in the same solution, provided they have sufficiently different fluorescent such as 10^{-9} g/cm³ could be detected. For example, this method is quite useful in the analysis of drugs, pesticides and atmospheric pollutants which are present in trace amounts. Studies based on fluorescence and phosphorescence provides important data on the properties excited states such as lifetime, energy and electronic configuration.

On the commercial side, fluorescent lamp is one of the applications of the phenomenon of fluorescence. A fluorescent lamp consists of a glass tube with

- a small amount of mercury,
- two electrodes, and
- a coating of phosphor.

A phosphor is a solid substance which emits fluorescent light when excited ultraviolet radiation. The electrodes initiate an electric arc which helps in vaporizing and exciting mercury atoms. The excited mercury atoms emit ultraviolet radiation. The phosphor, being excited by ultraviolet radiation, emits fluorescence.

Some other commercial applications of fluorescence are given below:

- Optical brighteners are added to detergents to give extra-brightness to the clothes. Optical brighteners fluoresce in sunlight.
- Fluorescent paints are manufactured using suitable additives.
- TV screens of different colours are produced using phosphors.

Although molecules in the excited a singlet and triple states could show interesting chemical behaviour, their studies are included in CHM 437 (Organic

Reaction Mechanism). In the next section, we shall discuss photosensitisation; this discussion could make you understand how a substance gains photochemical activity in the presence of another photochemically excited substance. Based on the materials of this section, answer the following SAQ.

SELF ASSESSMENT EXERCISE 5

State two differences between fluorescence and phosphorescence.

.....

5.6 Photosensitisation

Photosensitization is the process of exciting a molecule by energy transfer from an excited molecule. In this process, a donor molecule (D) absorbs a quantum light and forms an excited molecule (D^{*}). The excited donor molecule then transfers its excitation energy to an acceptor molecule (A) in the ground state in order to excite it. This can be explained using the following reaction sequence:


- Light absorption : $D + h\nu \rightarrow D^*$
- Sensitisation : $D^* + A \rightarrow A^* + D$


The donor molecule is called sensitizer. In Sec. 3.6, we mentioned about such energy transfer by the excited molecule. The excited acceptor molecule A^{*} could take part either in chemical reactions or in physical processes which we shall discuss now.

5.61 Photosensitised Chemical Reactions

If the excited molecule A^{*} has obtained sufficient energy, it will get dissociated and start a chemical reaction. The advantage in photosensitized dissociation of a molecule is that it is enough to transfer energy equivalent to its bond enthalpy to dissociate it. It is so since the photosensitized molecule gets dissociated in the ground state. This is in contrast to direct photochemical decomposition for which much higher energy is required due to the necessity of exciting a molecule to upper electronic state (see Sec. 3:4). Thus the energy required for photosensitized dissociation of a molecule is less than that of photochemical dissociation.


Let us consider an example. When irradiated with 253.7 nm light, hydrogen and oxygen react in presence of mercury vapour but not in its absence. Mercury vapour acts as a sensitizer. The reaction sequence is given below:





The chain reaction continues further. The energy transferred to hydrogen molecule by excited mercury atom is equal to 472 kJ mol^{-1} (as per Eq. 3:4 where $\lambda = 253.7\text{nm}$). This energy is sufficient for thermal dissociation of hydrogen since its bond enthalpy is 436 kJ mol^{-1} (see Table 3.2). Note that excited mercury atom cannot directly dissociate oxygen molecule since its bond enthalpy is higher (497 kJ mol^{-1}). Further in the absence of mercury vapour, light of 253.7 nm (or 472 kJ mol^{-1} energy) cannot photolyse hydrogen or oxygen directly, since the energies needed for their photochemical dissociation are much higher (1420 kJ mol^{-1} and 682 kJ mol^{-1} respectively). Hence mercury vapour is essential as a sensitizer for $\text{H}_2\text{-O}_2$ photochemical reaction as $\lambda = 253.7 \text{ nm}$.

A well-known photosensitized reaction is photosynthesis. Chlorophyll (“chl”) and other plant pigments act as photosensitizers in the synthesis of starch from carbon dioxide and water. A simplified reaction sequence is as follows:



Note:

Chlorophyll is the name given for a group of compounds with minor variation in structure. Chlorophyll-a absorbs effectively in the red region of sunlight; the red light is in abundance in sunlight.

The reaction mechanism is very complex. Energy calculations show that apart from chlorophyll, there must be other coloured light-absorbing materials (pigments) which also provide energy required for the synthesis of starch.

Photosensitization is frequently used by chemists for preparing compounds which cannot be formed by thermal or direct irradiation methods.

5.62 Sensitized Fluorescence

Let us now discuss a physical process which proceeds through sensations. Thallium vapour does not give rise to fluorescence when directly irradiated with light of wavelength 253.7nm . But if mercury vapour is also included in the reaction vessel, thallium exhibits fluorescence. Mercury atoms get excited first and transfer energy to thallium atoms to excite them. The excited thallium atoms emit fluorescent as they go down to the ground level.

Using the ideas learnt in the section, answer the following SAQ

SELF ASSESSMENT EXERCISE 6

Oxalic acid is not decomposed directly by light of wavelength, 335 nm. Irradiation in presence of uranyl (UO_2^+) ion results in the decomposition of oxalic acid to carbon monoxide and carbon dioxide.

- i) How do you explain this reaction?
- ii) Suggest a reaction sequence.

.....
.....
.....

(Note: the above procedure was used in the earlier chemical actinometrical method of determining the light intensity. Oxalic acid solution of known concentration was irradiated in presence of uranyl ion and the reacted oxalic acid was estimated using potassium permanganate solution. From the amount of oxalic acid reacted, the intensity of light was calculated using I for a particular wavelength.)

5.7 Applications Of Photochemistry

We have earlier indicated the applications of the study of fluorescence, phosphorescence, flash photolysis and photosensitisation. Apart from the above, a few are worth mentioning.

Synthetic organic chemists have increasingly started using photochemical methods for synthesis due to greater efficiency and selectively as compared to dark reaction. Photochemistry offers a method of conducting reactions which are not possible thermodynamically.

In the analysis of pollutants, photochemistry plays an important role. For example, photochemical studies have indicated how ozone layer is affected by chlorofluorocarbons (Freon) used as refrigerants, solvents and spray-propellants. The simplest Freon is CF_2Cl_2 . Freon is chemically inert and remains as such for years. But when it stratosphere (10 km to 50 km above the earth's surface), freon decomposes and gives out free chlorine atoms. These chlorine atoms can react with ozone decomposing it. This can cause depletion on zone layer. It is a matter of serious concern since ozone layers protects our planet from low wavelength portion of sun's rays (290 nm -320nm wavelength). Irradiation with such high energy radiation could product skin cancer. As a result of photochemical studies, alternatives are tried for Freon.


The present energy crisis has compelled the scientists to look for alternatives. Solar energy, if properly utilized through suitable photochemical reactions, could

offer a solution for this energy problem. In this connection we now discuss the functioning of a photogalvanic cell which is help for solar energy conversion through chemical reactions.

Notes:

Three common types of cells used for converting light into electricity are given below:

- i) Photoelectric cells or photo cells convert light into electricity using photosensitive cathode. (see Sec. 3.3)
- ii) Photovoltaic cells have two dissimilar silicon (or germanium) crystals in close contact. Irradiation of light causes flow of electrons from one crystal to other. Solar cells used in calculators are photovoltaic cells.
- iii) Photogalvanic cells convert light into electricity through chemical reactions.




Tris – (2,2' – bipyridine) ruthenium (II) ion

5.71 Photogalvanic Cell

A reversible cell which converts light into electricity through an intermediate chemical reaction is called photogalvanic cell. It undergoes cyclic charging and recharging process as explained below:

5.72 Charging in presence of light

Two substances undergo oxidation – reduction in presence of light



In this OA_1 and RA_1 refer to oxidized and reduced form of substance 1, while OA_2 and RA_2 to those of substance 2. In this process, light is used in conducting a chemical reaction. This photochemical reaction is thermodynamically non-spontaneous ($\Delta G > 0$). By the above photochemical process, the cell is charged.

5.73 Discharging in the absence of light

In the absence of light, the reaction reverses spontaneously ($\Delta G < 0$).



The chemical energy gained during charging is converted into electrical energy and the cell is discharged.

An example of such a system is given below:





Fig. 3.6: Photogalvanic cell




In this system, tris- (2,2'-bipyridine) ruthenium (II) – tris- (2,2'-bipyridine) ruthenium (III) and iron (II) – iron (III) are the two redox pairs. The platinum electrodes serve as electrical contacts (Fig. 3.6).

In the next section, we shall discuss chemiluminescence, which in a way, is the reverse of photochemical reaction.

Note:


Photochromism is the light induced reversible colour change. The colour change is brought about by reversible reactions. Photochromic sunglasses darken in sunlight and protect the eyes from the excess light. The sunglasses contain Ag^+ and react reversibly as follows:

In presence of sunlight:



Silver formed is responsible for the darkening of the glasses.

In the absence of sunlight:



SELF ASSESSMENT EXERCISE 7

Suggest a basic difference between galvanic cells (discussed in CHM 112 and photogalvanic cells

.....
.....
.....

5.74 Chemiluminescence

Chemiluminescence is the emission of light as a result of chemical reaction at room temperature. It must be clearly understood that chemiluminescence is not due to any photophysical process like fluorescence or phosphorescence. Some examples of chemiluminescence are given below:

- Glow of phosphorus is due to slow oxidation; it is not due to phosphorescence as name suggests.
- Oxidation of Grignard compounds by air or oxygen results in greenish-blue luminescence
- *Will-o-the-wisp* (mistaken as light produced by evil spirits) is the glow caused by the oxidation of decaying wood in marshy places.
- Emission of light by firefly is due to oxidation of some proteins in its body (this is also called bioluminescence).

5.8 SUMMARY

In this unit, we have discussed the physical and chemical processes accompanying light absorption.

- The laws of photochemistry have been stated and explained.
- Quantum yield has been defined and the method of determining it has been explained.
- The variation in energy requirements for photochemical and thermal dissociation has been discussed.
- The rate expressions have been derived from some photochemical reactions.

- The photophysical processes have been explained. The applications of fluorescence and phosphorescence have been stated.
- Photosensitization has been explained with examples.
- Chemiluminescence has been defined.

5.9 TUTOR MARKED ASSIGNMENT

- 1) State two differences between a chemical reaction and photochemical reaction.
- 2) If 10% of the energy of a 100 W incandescent bulb generates visible light having average wavelength 600nm, how many quanta of light are emitted in 10 minutes?

$$(1 \text{ W} = 1 \text{ J s}^{-1})$$

(Hint: a tungsten filament bulb is an example of incandescent bulb. Incandescence is light emission due to intense heat.)

- 3) The extinction coefficient ((ϵ)value of a substance is $4.66 \text{ m}^2 \text{ mol}^{-1}$. calculate the concentration of its solution (in molarity) which has $\frac{I}{I_0} = 0.2$. the thickness of the cell is 1.00 cm.
- 4) At 478nm, hydrogen and chlorine combine to give hydrogen chloride with a quantum efficiency of 1.00×10^6 . If the absorbed intensity is $2.50 \times 10^{-3} \text{ J s}^{-1}$, calculate the amount of hydrogen chloride formed in 10 minutes.

(Hints: The unit for amount is mole.)

- 5) State some of the applications of fluorescence and phosphorescence studies.
- 6) In the photochemical reaction between hydrogen and bromine at 511 nm wavelength, the first step is the photolysis of bromine but not of hydrogen, Why?
(Hints: Use table 2.2).
- 7) Fill in the blanks:
 - i) Light emission through $S_1 \rightarrow S_0$ transition is called.....
 - ii) Light emission through $T_1 \rightarrow S_0$ transition is called.....
 - iii) The energy required for photolysis of bromine is... than its bond enthalpy.
- 8) Suggest a condition that should be fulfilled in order to photolysis a molecule using a sensitizer.

5.10 ANSWERS TO SELF-ASSESSMENT QUESTIONS

$$1) \quad \text{Energy absorbed per photon} = \frac{6.626 \times 10^{-34} \times 3 \times 10^8}{250 \times 10^{-19}} \text{ joule photon}^{-1}$$

$$= 7.95 \times 10^{19} \text{ joule photon}^{-1}$$

Assume that the reaction is conducted for t seconds.

$$\text{Hence, } I_a = \frac{\text{Intensity in joule second}^{-1}}{7.95 \times 10^{19} \text{ joule photon}^{-1}}$$

$$= \frac{4.18 \times 10^3 \text{ joule} / t \text{ second}}{7.95 \times 10^{19} \text{ joule photon}^{-1}}$$

$$= \frac{5.26 \times 10^{21}}{t} \text{ photon second}^{-1}$$

$$\text{Amount of HI decomposed in 1 second} = \frac{1.85 \times 10^{-2} \text{ mol}}{t \text{ second}}$$

$$\frac{-dN_{\text{HI}}}{dt} = \frac{1.8510^{-2}}{t} \times 6.022 \times 10^{23} \text{ molecule second}^{-1}$$

$$= \frac{1.114 \times 10^{22}}{t} \text{ molecule second}^{-1}$$

$$\text{Using Eq. 3.3} = \frac{-dN_{\text{HI}}/dt}{I_a} = \frac{1.114 \times 10^{22} / t}{5.26 \times 10^{21} / t}$$

$$= 2.12$$

- 2) Oxygen molecules could quench the excited species. Hence in presence of oxygen, f_{HCl} is low.
- 3) For HI, the difference between photochemical dissociation energy and bond enthalpy is 68 kJ mol^{-1} and the excited hydrogen atom carries this excess energy.

4) i)
$$\frac{d[\text{CH}_3\text{CO}]}{dt} = 0 = k_2 [\text{CH}_3] [\text{CH}_3\text{CHO}] - k_3 [\text{CH}_3\text{CO}]$$

ii)
$$\frac{d[\text{CH}_3]}{dt} = 0 = I_a - k_2 [\text{CH}_3] [\text{CH}_3\text{CHO}] + k_3 [\text{CH}_3\text{CO}] - 2k_4 [\text{CH}_3]^2$$

iii)
$$\frac{d[\text{CH}_3\text{CO}]}{dt} + \frac{d[\text{CH}_3]}{dt} = 0 = I_a - 2k_4 [\text{CH}_3]^2$$

iv)
$$[\text{CH}_3] = \frac{I_a}{2k_4}$$

v)
$$\frac{d[\text{CO}]}{dt} = k_3 [\text{CH}_3\text{CO}]$$



vi)
$$\frac{d[\text{CO}]}{dt} = k_2 [\text{CH}_3] [\text{CH}_3\text{CHO}]$$

$$= k_2 \frac{I_a}{2k_4} [\text{CH}_3\text{CHO}]$$

- 5) The light emission during fluorescence is due to $S_1 \rightarrow S_0$ transition, fluorescence, being due to an allowed transition, is quite fast and stops as soon as exciting source is removed.

The light emission during phosphorescence is due to $T_1 \rightarrow S_0$ transition. Since this transition is forbidden, it is slow and it persists even if the exciting source is removed.

- 6) i) UO_2^+ ion acts as a photosensitiser and facilitates the decomposition of oxalic acid.



- 7) In a galvanic cell, the potential difference is due to conversion of chemical energy into electrical energy.

In a photogalvanic cell, light energy is converted into chemical energy, which is then, converted into electrical energy.

5.11 REFERENCES/FURTHER READINGS

Physical chemistry Joseph H Niggle 3rd edition Harper Collins college publisher 1936

Introduction to physical chemistry Mark ladd 3rd edition Cambridge university press 1999

Physical chemistry Atkins PW 3rd edition oxford university press 1986

Advanced chemistry (physical and industrial) Philip Mathew Cambridge university press 2003

Principles of Physical chemistry, Samuel H. Maron and Carl F. Prutton, Oxford and IBH Publishing Co., New Delhi, 4th ed., 1985. 1997.

Physical Chemistry, Gilbert W. Castellan, Narosa Publishing House, Addison-Wesley/Narosa (Indian student Edition), Delhi 3rd ed., 1983.

Physical Chemistry, P.C. Rakshit, Sarat Book House, Calcutta, 5th ed., 1988.

Physical Chemistry – Principles and Problems, D.V.S Jain and S.P. Jauhar, Tata McGraw-Hill Pub. Company Ltd., New Delhi, 1988.

APPENDIX I

Use of Eq. 2.66 in Calculating Total Collision Frequency

To understand the derivation of Eq. 2.79 for the total collision frequency among the molecules of two gases, X and Y, it is better to recapitulate the derivation of Eq. 2.52 given in Sec. 2.11 of Unit 2. Using a different procedure, Eq. 2.52 was derived for calculating the total collision frequency among the molecules of a Single gas.

Eq. 2.66 can be used for deriving the total collision frequency among the molecules of a single or two different gases. Now, let us see how Eq. 2.66 could be used to derive the total collision frequency among the molecules of a single gas.

Collision diameter of a single gas = σ

$$\left\{ \begin{array}{l} \text{The reduced mass of a system} \\ \text{of two molecules of the same} \\ \text{gas which undergo collision} \end{array} \right\} = \mu = \frac{m^2}{2m} = \frac{m}{2}$$

Mass of each molecule (m) is same and $m_x = m_y = m$ in Eq. 2.70

Mass of one molecule of a gas (m)

$$= \frac{\text{Molar mass } (M_m)}{\text{Avogadro constant } (N_A)}$$

And $k_b N_A = R$

Note that while dealing with relative motion of molecules, we must use reduced mass of a system of two molecules put not the mass of a molecule. Using the μ value given above in Eq. 2.68

$$\begin{aligned} \text{average relative speed} &= \left(\frac{8k_b T}{\pi m/2} \right)^{1/2} 2^{1/2} \left(\frac{8k_b T}{\pi m} \right)^{1/2} \\ \text{of two molecules of a gas} &= 2^{1/2} \left(\frac{8k_b T N_A}{\pi m N_A} \right)^{1/2} = 2^{1/2} \left(\frac{8RT}{\pi M_m} \right)^{1/2} \end{aligned}$$

$$\begin{aligned} \text{i.e., Average relative speed of} & \quad \left(\frac{8RT}{\pi M_m} \right)^{1/2} \\ \text{two molecules of gas} & \quad \left(\frac{\pi M_m}{8RT} \right)^{1/2} \\ & = 2^{1/2} \bar{u} \end{aligned}$$

Mass of one molecule of a gas (m) $\frac{\text{Molar mass (Mm)}}{\text{Avogadro constant (N}_A\text{)}}$ and $k_b N_A = R$
--

Number density of the gas = n_0

Now, let us come to the correction factor required while calculating the total collision frequency among the molecules of a single gas. Suppose there are N molecules in a gas with labels, 1, 2, 3, 4 ... N . To Calculate the total number of collisions, we use the following method:

First, imagine that molecule 1 is colliding with each of the molecules 2; 3, 4.....
 N and count the number of collisions

Then, assume that molecule 2 is colliding with each of the molecules, 1, 3, 4 ... N
 and count the number of collisions

Continue the above procedure till the N th molecule colliding with each of the
 molecules, 1, 2, 3 ... $(N - 1)$ and count the number of collisions in each case.

Add the number of collisions obtained in each case. In this procedure, each
 collision would have been counted twice. For example the collision between
 molecules 1 and 2 is counted twice as if, molecule 1 is colliding with molecule 2
 and, molecule 2 is colliding with molecule 1. Hence the correction factor, $1/2$,
 must be included in Eq. 2.68 while calculating the total collision frequency among
 the molecules of a gas, so that each collision is counted only once;

i.e., correction factor = $1/2$

Substituting the various factors in Eq. 2.66

$$\left. \begin{array}{l} \text{Total collision frequency} \\ \text{among the molecules of a gas} \\ \text{(denoted as } Z_{11} \text{ in Sec. 2.11)} \end{array} \right\} = \pi \sigma^2 \cdot 2^{1/2} \cdot u \cdot n_0 \cdot n_0 \cdot 1/2$$

$$= \frac{1}{\sqrt{2}} \pi \sigma^2 u n_0^2$$

While calculating the total collision frequency among the molecules of two
 different gases such as X and Y discussed in this section, there is no necessity for
 the corrector factor. We add up the number of collisions that each molecule of X
 will make with each molecules of Y as follows:

Let the molecules of gas X have labels,

$X_1, X_2, X_3 \dots X_N$

Let the molecules of gas Y have labels,

$Y_1, Y_2, Y_3 \dots Y_N$

Count the number of collisions as indicated below:

Let the molecule X_1 collide with each of the molecules, $Y_1, Y_2, Y_3 \dots Y_N$ and count the number of collisions.

Repeat the process and count the number of collisions that X_2 makes with $Y_1, Y_2, Y_3 \dots Y_N$. Continue the process and calculate the number of collisions upto the molecule X_N colliding with each of the molecules, $Y_1, Y_2, Y_3 \dots Y_N$

Add the number of collisions in each case.

In this process, each collision between a molecule of X and a molecule of Y is counted once only. Hence there is no necessity for correction factor while calculating the total collision frequency among the molecules of different gases.

In this case we have not **counted**

- i) the collision **among** the molecules of X only or
- ii) the collisions **among** the molecules of Y only.

APPENDIX II

Derivation of Eq. 2.97: $\Delta H^\ddagger = E_a - 2RT$

According to Eq. 2.59, $\ln k = \ln A - E_a/RT$. Differentiating with respect to temperature and assuming A and E_a to be independent of temperature,

$$\frac{d \ln k}{dT} = \frac{E_a}{RT^2}$$

or
$$E_a = RT^2 \frac{d \ln k}{dT}$$

This equation can be represented in the following form under constant volume conditions:

$$E_a = RT^2 \left(\frac{\partial \ln k}{\partial T} \right)_v \dots(A.1)$$

The change in concentration of a species is related to

- the change in the amount of the species and
- the change in the volume of the system.

If volume of the system is kept constant, the change in concentration of the species could be directly related to the change in the amount of species.

Keeping this in view, many reaction rates are measured under constant volume conditions. Reactions taking place in closed vessels are examples of this type.

$$E_a = RT^2 \left(\frac{\partial k/k}{\partial T} \right)_v \frac{RT^2}{k} = \left(\frac{\partial k}{\partial T} \right)_v \quad \dots(\text{A.2})$$

Rewriting Eq. 2.91

$$K = \frac{RT}{p^0} \frac{k_b T}{h} K_p$$

$$K = V^0 \frac{k_b T}{h} K_p \quad \dots(\text{A.3})$$

where $\frac{RT}{p^0} = V^0 = \text{standard volume}$

$$\left(\frac{\partial k}{\partial T} \right)_v = \frac{V^0 k_b}{h} \left(\frac{\partial(TK_p)}{\partial T} \right)_v$$

$$\left(\frac{\partial k}{\partial T} \right)_v = \frac{V^0 k_b}{h} \left[K_p + T \left(\frac{\partial K_p}{\partial T} \right)_v \right] \quad \dots(\text{A.4})$$

Using Eqs. A.3 and A.4

$$\frac{1}{k} \left(\frac{\partial k}{\partial T} \right)_v = \frac{h}{V k_b T K_p} \frac{V^0 k_b}{h} \left[K_p + T \left(\frac{\partial K_p}{\partial T} \right)_v \right]$$

$$= \frac{1}{TK_p} \left[K_p + T \left(\frac{\partial K_p}{\partial T} \right)_v \right] \quad \dots (\text{A.5})$$

$$\frac{RT^2}{k} \left(\frac{\partial k}{\partial T} \right)_v = \frac{RT^2}{TK_p} \left[K_p + T \left(\frac{\partial K_p}{\partial T} \right)_v \right]$$

$$= \frac{RT}{K_p} \left[K_p + T \left(\frac{\partial K_p}{\partial T} \right)_v \right]$$

$$\frac{RT^2}{k} \left(\frac{\partial k}{\partial T} \right)_v = RT + RT^2 \left(\frac{\partial \ln k_p}{\partial T} \right)_v \quad \dots(\text{A.6})$$

Using (A.2) and (A.6)

$$E_a = RT + RT^2 \left(\frac{\partial \ln k_p}{\partial T} \right)_v \quad \dots (\text{A.7})$$

$$-d \ln k_p = \frac{\Delta H^\circ}{RT^2} \dots\dots\dots (2.102)$$

Under constant volume conditions, takes the form.

$$RT^2 \left(\frac{\partial \ln k_p}{\partial T} \right)_v = \Delta U^\ddagger \quad \dots (\text{A.8})$$

where ΔU^\ddagger is the molar activation internal energy.

From Eqs. A.7 and A.8,

$$E_a = RT + \Delta U^\ddagger \quad \dots (\text{A.9})$$

Applying Eq. 4.15 pf CHE 112 for the formation of the activated complex,

$$\Delta H^\ddagger = \Delta U^\ddagger + \Delta n_g RT$$

Δn_g = Difference in the number of moles between the gaseous reactants and the gaseous products.

$$= -1 \text{ (for the formation of the activated complex as per Eq. 2.89)} \quad \dots(\text{A.10})$$

Hence $\Delta H^\ddagger = \Delta U^\ddagger - RT$

Using Eqs. A.9 and A.10,

$$E_a = \Delta H^\ddagger = \Delta U^\ddagger + 2RT$$

or $\Delta H^\ddagger = E_a - 2RT$

The last one is Eq. 2.97

Eq. 18.160 written below specifying the physical state $X(g) + Y \rightarrow A^\ddagger(g)$